

A General Model of Opinion Dynamics with Tunable Sensitivity

Anastasia Bizyaeva, Alessio Franci, and Naomi Ehrich Leonard

Abstract—We introduce a general model of continuous-time opinion dynamics for an arbitrary number of agents that communicate over a network and form real-valued opinions about an arbitrary number of options. Drawing inspiration from models in biology, physics, and social psychology, we apply a sigmoidal saturating function to inter-agent and intra-agent exchanges of opinions. The saturating function is the only nonlinearity in the model, yet we prove how it yields rapid and reliable formation of consensus, dissensus, and opinion cascades as a function of just a few parameters. We further show how the network opinion dynamics exhibit both robustness to disturbance and ultrasensitivity to inputs. We design feedback dynamics for system parameters that enable active tuning of implicit thresholds in opinion formation for sensitivity to inputs, robustness to changes in input, opinion cascades, and flexible transitions between consensus and dissensus. The general model can be used for systematic control design in a range of engineering problems including network systems, multi-robot coordination, task allocation, and decision making for spatial navigation. It can also be used for systematic examination of questions in biology and social science ranging from cognitive control and networks in the brain to resilience in collective animal behavior to changing environmental conditions to information spreading and political polarization in social networks.

Index Terms—opinion dynamics, decision making, networks, consensus, dissensus, cascade dynamics, sensitivity, robustness, bifurcations, nonlinear dynamics, control theory

I. INTRODUCTION

Opinion dynamics of networked agents are the subject of long-standing interdisciplinary interest. Agent-based models are often created to study local mechanisms that drive consensus formation and opinion clustering in a group. Such models are explored, for example, in studies of collective behavior in animal groups, of voting patterns, and of political polarization in human social networks. In engineering, they are fundamental to designing distributed coordination of autonomous agents and dynamic allocation of tasks within a network.

These models are typically used to investigate parameter regimes and network structures for which opinions in a group

converge over time to a desired configuration. However natural collectives have a remarkable ability not only to converge to a collective behavior, but also to rapidly switch between different collective behaviors in response to relevant changes in the surrounding environment. Understanding these temporal dynamics and how this ultrasensitive ability of natural groups to pick out meaningful information comes about is pivotal to our ability to design adaptable yet robust control laws for robotic teams and other networked engineered collectives.

This motivates us to pose the following questions, which we explore throughout the paper. How can a network of decision makers come rapidly and reliably to a meaningful configuration of opinions on multiple options, including consensus and dissensus, in response to internal biases or external stimuli? How can a network be controlled to do so with tunable sensitivity and robustness, distinguishing between meaningful signal and disturbance? How can a network reliably transition from one configuration of opinions to another in response to meaningful change? To investigate these questions, we present a new general agent-based dynamic model of the opinion formation process that is maximally rich in the behaviors it exhibits yet tractable to analysis by virtue of the small number of parameters needed to generate the full range of behaviors.

Classical models of opinion formation include [1]–[5]. In those models, agents’ opinions are represented by scalar values and linear *averaging mechanisms* rule agent interactions, commonly represented by a network graph with real-valued edge weights. Prominent nonlinear variations on averaging models include “bounded confidence” models, which assume that the opinion of a social agent is primarily influenced by its like-minded neighbors [6]–[9]. Consensus protocols on nonlinear manifolds are derived in [10], [11] by incorporating projections into averaging models. An extension of linear consensus protocols to networks with saturated inputs is considered in [12], [13].

In [14]–[18], nonlinear models are derived and studied in which agents with scalar opinions respond to a saturating function of the opinions of others, much like in a Hopfield neural network. In [14], the functional form represents *biased assimilation* from social psychology. The use of a saturating function has inspired the present development and the models provide a precursor to the general model defined here. Closely related to these are nonlinear models that leverage coupled oscillator dynamics [19], [20], biologically inspired mean-field models [21], or the Ising model for spin systems [22], [23].

To capture formation of opinions about multiple topics, extensions to classical models with vector-valued opinion states are proposed in [24]–[30]. Although importance of

This research has been supported in part by NSF grant CMMI-1635056, ONR grant N00014-19-1-2556, ARO grant W911NF-18-1-0325, DGAPA-UNAM PAPIIT grant IN102420, and Conacyt grant A1-S-10610. This material is also based upon work supported by the National Science Foundation Graduate Research Fellowship under Grant No. (NSF grant number).

A. Bizyaeva and N.E. Leonard are with the Department of Mechanical and Aerospace Engineering, Princeton University, Princeton, NJ, 08544 USA; e-mail: bizyaeva@princeton.edu, naomi@princeton.edu.

A. Franci is with the National Autonomous University of Mexico, 04510 Mexico City, Mexico. e-mail: afranci@ciencias.unam.mx

Any opinion, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation

logical interdependence between different topics is accounted for in [24], [26], [31], these models restrict each opinion to a binary choice, and topics are not typically assumed to be mutually exclusive. Conditions for consensus, clustering, and fragmentation of opinions are derived in [31].

In almost all of this literature, with [14], [18] being an exception, clustered or fragmented configurations of opinions typically arise only when an asymmetry or heterogeneity is imposed. For example, the emergence of opinion clusters is tied to special conditions on connectivity of a network in [3], [5], [18], [30]. In bounded confidence models, clustering of opinions is usually related to values of the opinion threshold, as in [29]. This implicitly creates a heterogeneous network structure among agents, since agents with sufficiently different opinions delete their communication link. Fragmentation or clustering of opinions have also been connected to heterogeneous biases or external information among agents [32].

In contrast to these findings, model-independent theory recently developed in [33] shows that opinion dynamics on an all-to-all network of homogeneous unbiased agents evaluating an arbitrary number of options can give rise to a fragmented global state. In this state, which we refer to as a type of group *dissensus*, the group is neutral about every option on average while individual agents are opinionated. Emergence of dissensus on a network is shown to be a *likely* outcome of opinion formation, alongside consensus, when the opinion formation process is nonlinear. This inherent coexistence of consensus and dissensus not tied to asymmetry or heterogeneity is a key feature of the proposed general model.

Further, the theory in [33] predicts coexistence and simultaneous stability of many different consensus and/or dissensus equilibria. This multistability can be understood to be fundamentally related to the type of tunably sensitive behavior observed in natural groups, since it enables a group to easily transition from one open configuration to another. Bistability of two consensus outcomes and the resulting sensitivity of collective opinion formation is explored for two options in [15]–[17]. Recent work in [34] similarly explores the role of bistability in opinion dynamics of a single agent recursively navigating a number of options organized in a binary tree hierarchy. Other nonlinear models of opinion formation among multiple options include [23], [35]. However, existing models of multi-option opinion dynamics for multi-agent systems do not capture maximally generic multistability of group outcomes.

In this paper we present a novel nonlinear model of multi-option opinion dynamics that captures consensus and dissensus outcomes predicted in [33] as well as the all important multistability of distinct consensus and/or dissensus equilibria. The structure of the model is derived by introducing the mechanisms of signal processing and evidence accumulation in networks of neurons into the model-independent structure constructed in [33]. Neuronal dynamics are inherently nonlinear and play a fundamental role in the underlying mechanisms behind the formation of opinions and decision making in animals and humans. Using their structure is therefore a logical modeling choice, which, as we show, captures the most general properties of the opinion formation process. This framework

provides a means to generate testable hypotheses about opinion dynamics and decision-making processes across a wide range of natural groups, including human social networks and animal groups. Further, such formulation allows us to mimic and take advantage of robustness and ultrasensitivity typical of neural systems to design novel distributed control approaches for engineered collectives.

The following are the major contributions of our work. First, we introduce a new general model for the study of multi-agent, multi-option opinion dynamics. The model allows a network to be defined on intra-agent and inter-agent opinion exchanges and includes an inertia term, an attention parameter that weights social influence, and an input term that can represent, for example, external stimuli, bias, or persistent opinions. Second, we show that with the introduction of saturating functions on opinion exchanges, the general model inherits the richness of generic opinion-formation behaviors predicted by the model-independent theory developed in the companion paper [33]: consensus, dissensus, and flexible transitions between them, even the highly symmetric homogeneous case of equal agents and equally valuable options; robustness of the opinion formation process to parameter heterogeneity and uncertainty; a small number of ruling parameters and analytical tractability. Third, we show how the model specializes to a range of models in the literature and in particular how the necessity of structurally balanced networks for clustered opinion formation in linear models breaks down in the nonlinear setting. Fourth, we introduce a distributed feedback dynamics to the agent attention parameters. Fifth, we show how attention feedback design parameters allow tunable sensitivity of opinion formation to inputs and its robustness to changes in inputs, tunable opinion cascades, even in response to a single agent receiving an input, and tunable transitions between consensus and dissensus.

The paper is organized as follows. In Section II we introduce notation and definitions. We present the new general model in Section III and show conditions for when it reduces to dynamics of agent clusters. In Section IV we prove how consensus and dissensus equilibria and multistability of opinion formation outcomes emerge as bifurcations of the general model, we discuss and illustrate the robustness of the opinion formation process to small heterogeneity. We show how the general model specializes to models in the literature and prove implications in Section V. We propose feedback dynamics for the attention parameter in Section VI and prove and illustrate tunable sensitivity, robustness, cascades, and transitions between consensus and dissensus. We conclude in Section VII.

II. NOTATION AND DEFINITIONS

Consider a network of N_a agents forming opinions about N_o options. Let $x_{ij} \geq 0$ be the magnitude of the *absolute* opinion of agent i about option j and suppose that every agent has the same total voting capacity:

$$x_{i1} + \dots + x_{iN_o} = r, \quad r > 0. \quad (1)$$

Absolute opinion state of agent i is $\mathbf{X}_i = (x_{i1}, \dots, x_{iN_o}) \in \Delta$, the $(N_o - 1)$ -dimensional simplex, and *absolute opinion state*

of the system is $\mathbf{X} = (\mathbf{X}_1, \dots, \mathbf{X}_{N_a}) \in \mathcal{V} = \Delta \times \dots \times \Delta$. At the neutral point $\mathcal{O} = (\mathcal{O}_1, \dots, \mathcal{O}_{N_a}) \in \mathcal{V}$, each agent has the same opinion about each option: $\mathcal{O}_i = \left(\frac{r}{N_o}, \dots, \frac{r}{N_o}\right) \in \Delta$.

In this paper we design and analyze opinion dynamics on the linear space $V = T_{\mathcal{O}}\mathcal{V} \subset \mathbb{R}^{N_a N_o}$ (the tangent space to \mathcal{V} at \mathcal{O}), where $V = V_a \times \dots \times V_a$ and $V_a = \{(v_1, \dots, v_{N_o}) \in \mathbb{R}^{N_o} \mid \sum_{l=1}^{N_o} v_l = 0\}$. As long as V is forward-invariant for the dynamics and the dynamics are bounded, they can be mapped back to the opinion dynamics on simplex space \mathcal{V} with an affine coordinate change (proved in Section III for our model).

Let $\mathbf{Z}_i = (z_{i1}, \dots, z_{iN_o}) \in V_a$ be the *opinion state of agent i* and $\mathbf{Z} = (\mathbf{Z}_1, \dots, \mathbf{Z}_{N_a}) \in V$ the *opinion state of the system*. These represent *relative* opinions: z_{ij} is the opinion of agent i about option j *relative* to the other options:

$$z_{i1} + \dots + z_{iN_o} = 0. \quad (2)$$

The neutral point $\mathbf{X} = \mathcal{O} \in \mathcal{V}$ corresponds to $\mathbf{Z} = \mathbf{0} \in V$.

Agent i is *unopinionated* if its opinion state is close to the neutral point: $\|\mathbf{Z}_i\| \leq \vartheta$, for $\vartheta \geq 0$ small and $\|\cdot\| = \|\cdot\|_2$. When $\|\mathbf{Z}_i\| > \vartheta$, agent i is *opinionated*. Agent i *favours* option j when it is opinionated and $z_{ij} \geq z_{ip} - \vartheta$ for all $p \neq j$. Conversely, agent i *disfavours* option j if it is opinionated and $z_{ij} < z_{ip} - \vartheta$ for some p . An agent is *conflicted* among a set of options if it has near equal and favorable opinions about all options in the set relative to the options not in the set.

Two agents *agree* if both are opinionated and share the same qualitative opinion state (e.g. favoring option j or conflicted among a set of options). Two agents *disagree* if they have different qualitative opinion states. If all agents are opinionated and agree, then the group is in an *agreement state*. If the group is in agreement and $\|\mathbf{Z}_i - \mathbf{Z}_k\| \leq \vartheta$, for all i, k , then the group is in a *consensus state*. If at least one pair of agents disagrees, the group is in a *disagreement state*. If the group is in disagreement and the average agent is unopinionated, i.e., $\|\frac{1}{N_a} \sum_{i=1}^{N_a} \mathbf{Z}_i\| \leq \vartheta$, then the group is in a *dissensus state*. The group is in an *unopinionated state* if all agents are unopinionated.

The system opinion state-space decomposes as $V = W_c \oplus W_d$, where W_c is the *multi-option consensus space* defined as

$$W_c = \{(\mathbf{Z}_1, \dots, \mathbf{Z}_{N_a}) \mid \mathbf{Z}_i = \tilde{\mathbf{Z}} \in V_a, \forall i\}, \quad (3)$$

and W_d is the *multi-option dissensus space* defined as

$$W_d = \{(\mathbf{Z}_1, \dots, \mathbf{Z}_{N_a}) \mid \mathbf{Z}_1 + \dots + \mathbf{Z}_{N_a} = \mathbf{0}\}. \quad (4)$$

On the consensus space W_c , agents have identical opinions. On the dissensus space W_d , agent opinions are balanced over the options such that the average opinion of the group is neutral.

Symmetry and equivariance: Let Γ be a compact Lie group acting on \mathbb{R}^n . Consider a dynamical system $\dot{\mathbf{x}} = \mathbf{h}(\mathbf{x})$ where $\mathbf{x} \in \mathbb{R}^n$ and $\mathbf{h} : \mathbb{R}^n \rightarrow \mathbb{R}^n$. Then $\rho \in \Gamma$ is a *symmetry* of the system, equivalently \mathbf{h} is ρ -*equivariant*, if $\rho \mathbf{h}(\mathbf{x}) = \mathbf{h}(\rho \mathbf{x})$. If \mathbf{h} is ρ -equivariant for all $\rho \in \Gamma$, then \mathbf{h} is Γ -*equivariant* [36]. In other words Γ -equivariance means that elements of the symmetry group Γ send solutions to solutions.

The compact Lie group associated with permutation symmetries of n objects is the *symmetric group on n symbols* S_n , which is the set of all bijections of $\Omega_n := \{1, \dots, n\}$ to

itself (i.e., all permutations of ordered sets of n elements). *Maximally symmetric* opinion dynamics are $(S_{N_a} \times S_{N_o})$ -equivariant, where elements of S_{N_a} permute the N_a -element set of agents and elements of S_{N_o} permute the N_o -element set of options [33]. Thus, maximally symmetric opinion dynamics are unchanged under any permutation of agents or options.

A subgroup $\Gamma_n \subset S_n$ is *transitive* if the orbit $\Gamma_n(i) = \{\gamma(i), \gamma \in \Gamma_n\} = \Omega$, for some (and thus all) $i \in \Omega$. $(\Gamma_{N_a} \times \Gamma_{N_o})$ -equivariant opinion dynamics, with transitive Γ_{N_a} and Γ_{N_o} , are still *highly symmetric* since any pair of agents and any pair of options, while not necessarily interchangeable by arbitrary permutations, can be mapped into each other by the symmetry group action. For example, if $\Gamma_{N_a} = \mathbf{D}_{N_a}$, the (transitive) dihedral group of order N_a , symmetries correspond to N_a rotations and N_a reflections. Thus, \mathbf{D}_{N_a} -equivariant opinion dynamics are unchanged if agents are permuted by a rotation or a reflection, e.g., if agents communicate over a network defined by a cycle.

Highly symmetric cases serve as the *organizing centers* [37] for the most generic opinion dynamics. The study of organizing centers and their perturbations (known as *unfoldings*) is the subject of *singularity theory* [37]. The theory provides the means to systematically describe the finite set of likely and qualitatively distinct dynamical behaviors as a function of parameters in the highly symmetric cases, with and without small heterogeneous inputs, perturbations, and uncertainties.

The general model of opinion dynamics, presented next, specializes to the highly symmetric cases, where formation of a consensus state (near W_c) or a dissensus state (near W_d) are the likely dynamical behaviors. We can then study how consensus and dissensus are affected by inputs and heterogeneity, and how they can be controlled with feedback.

III. GENERAL MODEL OF OPINION DYNAMICS

The general model describes continuous-time opinion dynamics for an arbitrary number N_a of agents (each representing an individual or a subgroup) that communicate over a network and form opinions about an arbitrary number N_o of options. For subsets of agents with similar features, we show in Section III-C a reduction to dynamics on clusters of agents.

A key feature of the model is the application of a saturating function to exchanges of opinions among agents. Dynamics that evolve according to saturating interactions appear in Hopfield neural network models, which describe the emergence of associative memories [38]–[40], and in nonlinear leaky competing accumulator models, which relate decision making among alternatives to integration of noisy evidence in the brain [41], [42]. Likewise, in the discrete-time model of opinion formation for two options of [14], [18], a saturating Hill function is used to represent “biased assimilation” of social information, a well-studied concept from social psychology.

In the general model, the saturating function is the only nonlinearity. Yet, it is sufficient to yield consensus and dissensus as a function of just a few parameters, even in the homogeneous and maximally symmetric case. This is not the case for opinion dynamics with a linear averaging term (e.g., linear variations of the DeGroot model [1]), even allowing for antagonistic interconnections as in [5]; see Section V.




Fig. 1: Illustration of the four classes of interactions.

A. Model

The general model defines the rate of change of z_{ij} , agent i 's opinion about option j , as a function of three terms: an inertia term, a social term, and an input term.

The *inertia* term, parametrized by $d_{ij} > 0$, on its own drives z_{ij} exponentially in time to 0 (neutral opinion). In the general model, a larger d_{ij} implies a greater resistance of agent i to forming a non-neutral opinion about option j .

The *social* term is defined as the product of an *attention* parameter $u_i \geq 0$ and a saturating function of weighted sums of agent opinions that are available to agent i and influence its opinion of option j . The social term can also be interpreted as an *activation* term. The weights are encoded in adjacency tensor $A \in \mathbb{R}^{N_a \times N_a \times N_o \times N_o}$, such that z_{ij} is affected by z_{kl} only if $A_{ik}^{jl} \in \mathbb{R}$ is nonzero. The magnitude of A_{ik}^{jl} determines the strength of influence of agent k 's opinion about option l on agent i 's opinion about option j , and the sign of A_{ik}^{jl} determines whether this interaction is *excitatory* ($A_{ik}^{jl} > 0$) or *inhibitory* ($A_{ik}^{jl} < 0$). We reserve indices i, k to refer to agents and indices j, l to refer to options.

We distinguish four classes of interactions (see Fig. 1):

- 1) *Intra-agent, same-option coupling*: A_{ii}^{jj}
- 2) *Intra-agent, inter-option coupling*: $A_{ii}^{jl}, j \neq l$
- 3) *Inter-agent, same-option coupling*: $A_{ik}^{jj}, i \neq k$
- 4) *Inter-agent, inter-option coupling*: $A_{ik}^{jl}, i \neq k, j \neq l$.

Because options are mutually exclusive, agents i and k are *cooperative* if $\min\{A_{ik}^{jj}, A_{ki}^{jj}\} > \max\{A_{ik}^{jl}, A_{ki}^{jl}\}$, for all options $j \neq l$, i.e., the opinion of agent i (agent k) about option j is more strongly excited by the opinion of agent k (agent i) about the same option j than by its opinion about a distinct option l . Conversely, agents i and k are *competitive* if $\max\{A_{ik}^{jj}, A_{ki}^{jj}\} < \min\{A_{ik}^{jl}, A_{ki}^{jl}\}$.

The *attention* parameter $u_i \geq 0$ governs the strength of the social term relative to inertia and can have its own dynamics. We show in Section VI how u_i can be used as a dynamic feedback control parameter to tune sensitivity of the opinion dynamics to input. Attention u_i can model *urgency* in adopting a non-neutral opinion by having it grow with increasing time-urgency (election day approaching), or *spatial-urgency* (target getting closer). u_i can also be used to model *social effort*, *excitability*, or *susceptibility* of agent i to social influence.

The *input* $b_{ij} \in \mathbb{R}$ represents an input signal from the environment or a bias or predisposition that directly affects

agent i 's opinion of option j . For example, the input b_{ij} can be used to model the exogenous influence of agent i 's initial opinions, as in [2], where agents hold on to their initial opinions (sometimes called "stubborn" agents as in [43]).

The *general model of opinion dynamics* is given, for every agent $i = 1, \dots, N_a$ and every option $j \in 1, \dots, N_o$, by

$$\dot{z}_{ij} = F_{ij}(\mathbf{Z}) - \frac{1}{N_o} \sum_{l=1}^{N_o} F_{il}(\mathbf{Z}) \quad (5a)$$

$$F_{ij}(\mathbf{Z}) = -d_{ij}z_{ij} \quad (5b)$$

$$+ u_i \left(S_1 \left(\sum_{k=1}^{N_a} A_{ik}^{jj} z_{kj} \right) + \sum_{\substack{l=1 \\ l \neq j}}^{N_o} S_2 \left(\sum_{k=1}^{N_a} A_{ik}^{jl} z_{kl} \right) \right) + b_{ij}.$$

The drift $F_{ij} : V \rightarrow \mathbb{R}$, defined in (5b), is the sum of the three influences on agent i 's opinion of option j : inertia, social, and input. $S_q : \mathbb{R} \rightarrow [-k_{q1}, k_{q2}]$ with $k_{q1}, k_{q2} \in \mathbb{R}^{>0}$ for $q \in \{1, 2\}$ is a generic sigmoidal saturating function satisfying constraints $S_q(-x) \neq -S_q(x)$, $S_q(0) = 0$, $S'_q(0) = 1$, $S''_q(0) \neq 0$, $S'''_q(0) \neq 0$. S_1 saturates same-option interactions, and S_2 saturates inter-option interactions. S_1 and S_2 could be the same but are distinguished in (5) for the most general statement of the model. The rate of change of z_{ij} is defined in (5a) to ensure that z_{ij} remains agent i 's opinion of option j relative to its opinions of the other options. Subtracting the average drift over options in (5a) models the mutual exclusivity of options and makes state space V forward invariant for (5), as proved in Lemma III.3 below.

Let $\hat{b}_i = \frac{1}{N_o} \sum_{l=1}^{N_o} b_{il}$ be the *average input to agent i* and let $b_{ij}^\perp = b_{ij} - \hat{b}_i$ be the *relative input to agent i for option j* .

Lemma III.1. *The average input \hat{b}_i for each agent i has no effect on the emergent dynamics of the network.*

Proof. Define $G_{ij}(\mathbf{Z}) = F_{ij}(\mathbf{Z}) - b_{ij}$, $G'_{ij} = G_{ij} - \sum_{p=1}^{N_o} G_{ip}$. Then $\dot{z}_{ij} = G'_{ij}(\mathbf{Z}) + b_{ij} - \frac{1}{N_o} \sum_{p=1}^{N_o} b_{ip} = G_{ij}(\mathbf{Z}) + b_{ij}^\perp$. \square

Lemma III.1 implies that only relative inputs affect the opinion dynamics. Without relative inputs, the system (5) always has the neutral point as an equilibrium.

Lemma III.2. *$\mathbf{Z} = \mathbf{0}$ is an equilibrium for (5) if and only if there are no relative inputs, i.e., $b_{ij}^\perp = 0$ for all i and all j .*

When relative inputs are small, i.e., they do not dominate the dynamics, the formation of opinions in the general model (5) is governed by the balance between the inertia term, which resists opinion formation, and the social term, which promotes opinion formation. For illustrative purposes, consider the case in which $u_i = u \geq 0$ for all i . Then for u small, inertia dominates and the system behaves linearly. The opinions z_{ij} remain small and their relative magnitude is determined by the small inputs b_{ij}^\perp . For u large, the social term dominates and the system behaves nonlinearly. Opinions z_{ij} form that are much larger than, and potentially unrelated to, inputs b_{ij}^\perp , even for very small initial conditions. That opinions reach consensus or dissensus states for (and near) the highly symmetric cases is predicted by the model-independent theory of [33]. This theory also predicts that the opinion formation will be rapid

in general, and like a switch for $N_o \geq 3$. As we will show, the general model (5) has the great advantage that it exhibits the range of predicted behaviors with only a few parameters.

B. Well-Definedness of Model

We show that the general model (5) is well defined by showing in Lemma III.3 that V is forward invariant for (5) and in Theorem III.5 that solutions are bounded. This implies, as we show in Corollary III.5.1, that (5) can be mapped from V to the simplex product space \mathcal{V} , so that the general model (5) describes opinion dynamics of agents with equal voting capacity. The simplex product space \mathcal{V} is often associated with models of opinion dynamics, e.g., in [32], [44], [45].

Lemma III.3. V is forward invariant for (5).

Proof. For all $i = 1, \dots, N_a$, $\sum_{j=1}^{N_o} \dot{z}_{ij} = 0$, so if $z_{i1}(0) + \dots + z_{iN_o}(0) = 0$, $z_{i1}(t) + \dots + z_{iN_o}(t) = 0$ for all $t > 0$. \square

To show boundedness of solutions of (5), we first examine the case in which $u_i = 0$ for every agent i , and we prove global exponential stability of the corresponding unique equilibrium. When $u_i = 0$ for every i , the general model reduces to

$$\dot{z}_{ij} = - \left(d_{ij} z_{ij} - \frac{1}{N_o} \sum_{l=1}^{N_o} d_{il} z_{il} \right) + b_{ij}^\perp. \quad (6)$$

We rewrite (6) in vector form as $\dot{\mathbf{Z}} = D\mathbf{Z} + \mathbf{b}^\perp$ where $\mathbf{b}^\perp = (b_{11}^\perp, \dots, b_{1N_o}^\perp, \dots, b_{N_a 1}^\perp, \dots, b_{N_a N_o}^\perp)$ and $D = \text{diag}\{D_i\} \in \mathbb{R}^{N_a N_o \times N_a N_o}$ with $D_i \in \mathbb{R}^{N_o \times N_o}$ given by

$$D_i = \begin{pmatrix} -\frac{N_o-1}{N_o} d_{i1} & \frac{1}{N_o} d_{i2} & \dots & \frac{1}{N_o} d_{iN_o} \\ \frac{1}{N_o} d_{i1} & -\frac{N_o-1}{N_o} d_{i2} & \dots & \frac{1}{N_o} d_{iN_o} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{1}{N_o} d_{i1} & \frac{1}{N_o} d_{i2} & \dots & -\frac{N_o-1}{N_o} d_{iN_o} \end{pmatrix}. \quad (7)$$

D_i is the Laplacian of a weighted all-to-all graph.

Theorem III.4. Consider linear dynamics (6) which are the general dynamics (5) with $u_i = 0$ for all i .

- i) If, $b_{ij}^\perp = 0$ for all i, j , then the neutral equilibrium $\mathbf{Z} = \mathbf{0}$ is globally exponentially stable in V .
- ii) If for at least one pair i, j , $b_{ij}^\perp \neq 0$, then a nontrivial equilibrium state $\tilde{\mathbf{Z}}^*$, fully determined by \mathbf{b}^\perp , is globally exponentially stable in V .

Proof. By the Greshgorin circle theorem, every eigenvalue λ_{ij} of D lies within the closed disk in \mathbb{C} , centered at $-\frac{N_o-1}{N_o} d_{ij}$ with radius $R_{ij} = \frac{N_o-1}{N_o} d_{ij}$. Since $d_{ij} > 0$, $\text{Re}[\lambda_{ij}] < 0$ or $\lambda_{ij} = 0$, for every eigenvalue λ_{ij} . The zero eigenvalue has multiplicity N_a with eigenvectors $\mathbf{n}_i = (\mathbf{0}, \dots, \mathbf{d}_i, \dots, \mathbf{0})$, where $\mathbf{0} \in \mathbb{R}^{N_o}$ is the vector of all zeros and $\mathbf{d}_i = \left(\frac{1}{d_{i1}}, \frac{1}{d_{i2}}, \dots, \frac{1}{d_{iN_o}} \right)$ is in the i th position. Let $\mathcal{N} = \text{span}\{\mathbf{n}_i\}$. Then $V \cap \mathcal{N} = \{\mathbf{0}\}$ and $\dim V + \dim \mathcal{N} = N_a N_o$.

By Lemma III.3, range $D = V$. So, V is spanned by the eigenvectors of D corresponding to λ_{ij} with $\text{Re}[\lambda_{ij}] < 0$. Thus, trajectories of (6) with $b_{ij}^\perp = 0$, restricted to the invariant subspace V , decay exponentially to $\mathbf{Z} = \mathbf{0}$. The result ii) follows directly from i) with a change of coordinates. \square

We use Theorem III.4, to prove bounded solutions of (5).

Theorem III.5 (Boundedness). Let \bar{U} be a compact subset of \mathbb{R} . There exists $R > 0$ such that, for all $u, d_{ij}, A_{ik}^{jl}, b_{ij} \in \bar{U}$, $i, k = 1, \dots, N_a$, $j, l = 1, \dots, N_o$, the set

$$V \cap \{|z_{ij}| \leq R, i = 1, \dots, N_a, j = 1, \dots, N_o\}$$

is forward invariant for (5). This implies that the solutions $\mathbf{Z}(t)$ of the dynamics (5) are bounded for all time $t \geq 0$.

Proof. By boundedness of $S_p(\cdot)$, there exists $\tilde{R} > 0$ such that, for all $u_i, d_{ij}, A_{ik}^{jl}, b_{ij} \in \bar{U}$, $F_{ij}(\mathbf{Z}) = -d_{ij} z_{ij} + C_{ij}(\mathbf{Z})$, with $|C_{ij}(\mathbf{Z})| \leq \tilde{R}$. For all $\mathbf{Z} \in V$, it holds that

$$\begin{aligned} \frac{d}{dt} \frac{1}{2} \|\mathbf{Z}\|^2 &= \sum_{i=1}^{N_a} \sum_{j=1}^{N_o} z_{ij} \dot{z}_{ij} \\ &= \sum_{i=1}^{N_a} \sum_{j=1}^{N_o} z_{ij} \left(-d_{ij} z_{ij} + C_{ij}(\mathbf{Z}) + \frac{1}{N_o} \sum_{l=1}^{N_o} (d_{il} z_{il} - C_{il}(\mathbf{Z})) \right) \\ &= \mathbf{Z}^T D \mathbf{Z} + \sum_{i=1}^{N_a} \sum_{j=1}^{N_o} z_{ij} \left(C_{ij}(\mathbf{Z}) - \frac{1}{N_o} \sum_{l=1}^{N_o} C_{il}(\mathbf{Z}) \right) \\ &\leq \mathbf{Z}^T D \mathbf{Z} + N_a N_o \tilde{R} \|\mathbf{Z}\| \end{aligned}$$

where we have used $\sum_{j=1}^{N_o} z_{ij} = 0$ for all i . We compute

$$\begin{aligned} \mathbf{Z}^T D \mathbf{Z} &= \sum_{i=1}^{N_a} \sum_{j=1}^{N_o} (-d_{ij} z_{ij}^2) + \frac{1}{N_o} \sum_{i=1}^{N_a} \sum_{l=1}^{N_o} d_{il} z_{il} \left(\sum_{j=1}^{N_o} z_{ij} \right) \\ &= \sum_{i=1}^{N_a} \sum_{j=1}^{N_o} -d_{ij} z_{ij}^2 \leq -\min_{i,j} \{d_{ij}\} \|\mathbf{Z}\|^2. \end{aligned}$$

Then, for all $\|\mathbf{Z}\| \geq \frac{N_a N_o \tilde{R}}{\min_{i,j} \{d_{ij}\}}$,

$$\frac{d}{dt} \frac{1}{2} \|\mathbf{Z}\|^2 \leq -\|\mathbf{Z}\| \left(\min_{i,j} \{d_{ij}\} \|\mathbf{Z}\| - N_a N_o \tilde{R} \right) \leq 0.$$

The result follows by [46, Theorem 4.18]. \square

Corollary III.5.1. Mapping to the Simplex Product \mathcal{V} . Given a bounded set $\bar{U} \subset \mathbb{R}$, assume $u, d_{ij}, A_{ik}^{jl}, b_{ij} \in \bar{U}$, $i, k = 1, \dots, N_a$, $j, l = 1, \dots, N_o$. Let r be defined by (1). Then, the vector field of (5) can be mapped from the forward invariant region $V \cap \{|z_{ij}| \leq R, i = 1, \dots, N_a, j = 1, \dots, N_o\}$ to the product of simplex \mathcal{V} by the affine change of coordinates

$$\begin{aligned} L : V \cap \{|z_{ij}| \leq R, i = 1, \dots, N_a, j = 1, \dots, N_o\} &\rightarrow \mathcal{V} \\ \mathbf{Z} &\mapsto \frac{r}{N_o R} \mathbf{Z} + \frac{r}{N_o}. \end{aligned}$$

C. Clustering and Model Reduction

We prove sufficient conditions for the dynamics (5) of N_a agents and N_o options to reduce to the dynamics of N_c clusters and N_o options, and we derive the reduced dynamics. Each cluster $p = 1, \dots, N_c$ represents N_p of the N_a agents forming opinions as a unit and $\sum_{p=1}^{N_c} N_p = N_a$.

Theorem III.6 (Model Reduction with Opinion Clusters). Suppose there are N_c clusters with N_p agents in the p th cluster such that $\sum_{p=1}^{N_c} N_p = N_a$. Let \mathcal{I}_p be the set of indices for agents in the p th cluster. Assume for every $p = 1, \dots, N_c$:

- 1) $u_i = u_p$, $d_{ij} = d_{pj}$, $b_{ij} = \hat{b}_i + b_{pj}^\perp$ for $i \in \mathcal{I}_p$;
- 2) $A_{ii}^{jj} = \bar{B}_{pp}^{jj}$, $A_{ik}^{jj} = B_{pp}^{jj}$, $A_{ii}^{jl} = \bar{B}_{pp}^{jl}$, $A_{ik}^{jl} = B_{pp}^{jl}$ for $i, k \in \mathcal{I}_p$, and $i \neq k$;
- 3) $A_{ik}^{jj} = B_{ps}^{jj}$, $A_{ik}^{jl} = B_{ps}^{jl}$ for $i \in \mathcal{I}_p$, $k \in \mathcal{I}_s$, $s = 1, \dots, N_c$ and $s \neq p$,

with $d_{pj} > 0$, $u_p \geq 0$, and $B_{ps}^{jl}, b_{pj}^\perp \in \mathbb{R}$ for all $p, s = 1, \dots, N_c$, $j, l = 1, \dots, N_o$, $p \neq s$, $j \neq l$. Define bounded set $K_q \subset \mathbb{R}^{>0}$, $q = 1, 2$, as the image of the derivative of the saturating function S'_q of (5). If the following condition holds:

$$\sup_{\kappa_1 \in K_1, \kappa_2 \in K_2} \left\{ -\min_j \{d_{pj}\} + u_p \kappa_1 \max_j \{\bar{B}_{pp}^{jj} - B_{pp}^{jj}\} + u_p \kappa_2 \max_j \{\bar{B}_{pp}^{jl} - B_{pp}^{jl}\} \right\} < 0 \quad \forall p = 1, \dots, N_c, \quad (8)$$

then every trajectory of (5) converges exponentially in time to a $N_c(N_o - 1)$ -dimensional attracting manifold defined by

$$\mathcal{E} = \{\mathbf{Z} \in V \mid z_{ij} = z_{kj} \quad \forall i, k \in \mathcal{I}_p, p = 1, \dots, N_c\}. \quad (9)$$

The dynamics on \mathcal{E} reduce to (5) with N_c agents with opinion states $\hat{z}_{pj} = z_{ij}$ for any $i \in \mathcal{I}_p$, $p = 1, \dots, N_c$, with weights

$$\hat{A}_{pp}^{jj} = \bar{B}_{pp}^{jj} + (N_p - 1)B_{pp}^{jj}, \quad \hat{A}_{ps}^{jj} = N_s B_{ps}^{jj}, \quad (10a)$$

$$\hat{A}_{pp}^{jl} = \bar{B}_{pp}^{jl} + (N_p - 1)B_{pp}^{jl}, \quad \hat{A}_{ps}^{jl} = N_s B_{ps}^{jl}. \quad (10b)$$

Proof. The opinion dynamics (5) of agent $i \in \mathcal{I}_p$ is defined by the drift term

$$F_{ij}(\mathbf{Z}) = -d_{pj}z_{ij} + u_p \left(S_1 \left(\bar{B}_{pp}^{jj}z_{ij} + B_{pp}^{jj} \sum_{k \in \mathcal{I}_p} z_{kj} + \sum_{\substack{s \neq p \\ s=1}}^{N_c} \sum_{k \in \mathcal{I}_s} B_{ps}^{jj}z_{kj} \right) + \sum_{\substack{l \neq j \\ l=1}}^{N_o} S_2 \left(\bar{B}_{pp}^{jl}z_{il} + B_{pp}^{jl} \sum_{k \in \mathcal{I}_p} z_{kl} + \sum_{\substack{s \neq p \\ s=1}}^{N_c} \sum_{k \in \mathcal{I}_s} B_{ps}^{jl}z_{kl} \right) \right) + \hat{b}_i + b_{pj}^\perp. \quad (11)$$

Let $V(\mathbf{Z}) = \sum_{p=1}^{N_c} V_p(\mathbf{Z})$, $V_p(\mathbf{Z}) = \frac{1}{2} \sum_{i, k \in \mathcal{I}_p} \sum_{j=1}^{N_o} (z_{ij} - z_{kj})^2$. Recall that $F_{ij}(\mathbf{Z}) = -d_{ij}z_{ij} + C_{ij}(\mathbf{Z})$. Then

$$\begin{aligned} \dot{V}_p(\mathbf{Z}) &= - \sum_{i \in \mathcal{I}_p} \sum_{k \in \mathcal{I}_p} (\mathbf{Z}_i - \mathbf{Z}_k)^T D_p(\mathbf{Z}_i - \mathbf{Z}_k) \\ &\quad + \sum_{i \in \mathcal{I}_p} \sum_{k \in \mathcal{I}_p} \sum_{j=1}^{N_o} (z_{ij} - z_{kj})(C_{ij}(\mathbf{Z}) - C_{kj}(\mathbf{Z})) \\ &\quad - \frac{1}{N_o} \sum_{i \in \mathcal{I}_p} \sum_{k \in \mathcal{I}_p} \sum_{j=1}^{N_o} \sum_{l=1}^{N_o} (z_{ij} - z_{kj})(C_{il}(\mathbf{Z}) - C_{kl}(\mathbf{Z})). \end{aligned} \quad (12)$$

where D_p is from (7). The third term in (12) is zero because $\sum_{j=1}^{N_o} z_{ij} = 0$ on V . By the Mean Value Theorem, we write $C_{ij}(\mathbf{Z}) - C_{kj}(\mathbf{Z})$ in the second term as

$$u_p (\kappa_1 (\bar{B}_{pp}^{jj} - B_{pp}^{jj}) - \kappa_2 (\bar{B}_{pp}^{jl} - B_{pp}^{jl})) (z_{ij} - z_{kj})^2 \quad (13)$$

where $\kappa_1 \in K_1$ and $\kappa_2 \in K_2$. Then we find that

$$\begin{aligned} \dot{V}_p(\mathbf{Z}) &\leq \sup_{\kappa_1 \in K_1, \kappa_2 \in K_2} \left\{ -\min_j \{d_{pj}\} + u_p \kappa_1 \max_j \{\bar{B}_{pp}^{jj} - B_{pp}^{jj}\} \right. \\ &\quad \left. + u_p \kappa_2 \max_j \{\bar{B}_{pp}^{jl} - B_{pp}^{jl}\} \right\} 2V_p(\mathbf{Z}). \end{aligned} \quad (14)$$

When (8) is satisfied, using LaSalle's invariance principle [46, Theorem 4.4] every trajectory of (5) converges exponentially in time to the largest invariant set of $V(\mathbf{x}) = 0$, which is \mathcal{E} . Let $\hat{z}_{pj} = z_{ij}$ for any $i \in \mathcal{I}_p$. The dynamics (11) on \mathcal{E} reduce to (5) with $N_a = N_c$ and weights (10). \square

Whenever conditions of Theorem III.6 are met, the group of N_a agents will converge to a clustered group opinion state. This can happen for a broad class of interaction networks but also for all-to-all network and interaction weights with the same sign; see Section V for an illustration with two options.

IV. CONSENSUS AND DISSENSUS

In this section we show how consensus *and* dissensus emerge generically for the opinion dynamics (5) in maximally and highly symmetric cases. This justifies our claim that the general model (5) realizes the full richness of opinion formation behavior, as identified by the model-independent theory of [33]. It further sets the stage for investigating the influence on opinion formation of symmetry-breaking inputs, as we do in Section VI, and of asymmetries such as irregular network structure, as we do in ongoing work.

We show here how formation of consensus versus dissensus is determined by a small number of parameters that distinguish whether the deciding agents are cooperative or competitive. We further explain and illustrate how the emergence of consensus and dissensus is robust to small inputs and small heterogeneity, perturbation, and uncertainty in all parameters. Another important result for the maximally and highly symmetric cases is multi-stability of equilibria, e.g., stability of consensus in favor of each of the N_o options. In fact, there are conditions on parameters for which consensus and dissensus solutions *co-exist*, revealing the possibility of easy transition from consensus to dissensus or the reverse. We conclude the section with a generalization of the consensus and dissensus results in the case of clustered dynamics.

We first state necessary and sufficient conditions for $(S_{N_o} \times S_{N_a})$ -equivariance of (5), the maximally symmetric case.

Proposition IV.1. *Model (5) is $(S_{N_o} \times S_{N_a})$ -equivariant if and only if $b_{ij} = \hat{b} \in \mathbb{R}$, $d_{ij} = d > 0$, $u_i = u \geq 0$, $A_{ii}^{jj} = \alpha \in \mathbb{R}$, $A_{ii}^{jl} = \beta \in \mathbb{R}$, $A_{ik}^{jj} = \gamma \in \mathbb{R}$, $A_{ik}^{jl} = \delta \in \mathbb{R}$ for all $i, k = 1, \dots, N_a$, $k \neq i$, and all $j, l = 1, \dots, N_o$, $j \neq l$.*

Proof. The proof follows analogously to that of [33, Theorem 2.5] with the additional coefficient d_{ij} on the linear terms. \square

By Proposition IV.1 maximal symmetry corresponds to a network of identical agents with no relative input. Maximal symmetry also requires each of the four types of network interactions outlined in Section III-A to be all-to-all and homogeneous, with coupling weights $\alpha, \beta, \gamma, \delta$, respectively.

Remark IV.1. *More generally, for identical agents with no relative input, the symmetry group of the opinion dynamics is determined by the automorphism group of the multi-graph associated to adjacency tensor A . For instance, if the inter-agent coupling topologies, as determined by $[A_{ik}^{jj}]$ and $[A_{ik}^{jl}]$, $i \neq k$, $j \neq l$, are undirected rings, then the agent symmetry is D_{N_a} , the dihedral group of order N_a (proved below). If the*

inter-agent coupling topologies are directed rings, then the agent symmetry is Z_{N_a} , the cyclic group of order N_a .

We next illustrate the above remark and prove sufficient conditions for $(S_{N_o} \times D_{N_a})$ -equivariance of (5) with agent communication topology connected in an undirected cycle, a highly symmetric case.

Proposition IV.2. *Consider the model (5) with $b_{ij} = \hat{b} \in \mathbb{R}$, $d_{ij} = d > 0$, $u_i = u \geq 0$, $A_{ii}^{jj} = \alpha \in \mathbb{R}$, $A_{ii}^{jl} = \beta \in \mathbb{R}$, $A_{ik}^{jj} = \gamma \tilde{a}_{ik}$, $A_{ik}^{jl} = \delta \tilde{a}_{ik}$, with $\gamma, \delta \in \mathbb{R}$ and $\tilde{a}_{ik} \in \{0, 1\}$ for all $i, k = 1, \dots, N_a$, $i \neq k$, and for all $j, l = 1, \dots, N_o$, $j \neq l$. If $(\tilde{a}_{ik}) := \tilde{A} \in \mathbb{R}^{N_a \times N_a}$ is the adjacency matrix of an undirected cycle graph, then model is $(S_{N_o} \times D_{N_a})$ -equivariant. and the group acts transitively on V .*

Proof. To prove equivariance it is sufficient to show that the dynamics are equivariant under the action of the generators of the symmetry group $S_{N_o} \times D_{N_a}$.

Elements $\sigma \in S_{N_o}$ act on V by permuting the elements z_{ij} of each agent's opinion vector \mathbf{Z}_i . Recall that the generators of S_{N_o} are N_o transpositions σ_j where each σ_j swaps adjacent elements j and $j+1$ (or N_o and 1 when $j = N_o$). Let $\mathbf{F}_i(\mathbf{Z}) = (F_{i1}(\mathbf{Z}), \dots, F_{iN_o}(\mathbf{Z}))$ and observe that $\sigma_j \mathbf{F}_i(\mathbf{Z}) = (F_{i1}(\mathbf{Z}), \dots, F_{i(j+1)}(\mathbf{Z}), F_{ij}(\mathbf{Z}), \dots, F_{iN_o}(\mathbf{Z}))$. Furthermore observe that computing $\mathbf{F}_i(\sigma_j \mathbf{Z})$, only F_{ij} and $F_{i(j+1)}$ are changed, with

$$\begin{aligned} F_{ij}(\sigma_j \mathbf{Z}) &= -dz_{i(j+1)} \\ &+ u \left(S_1 (\alpha z_{i(j+1)} + \gamma z_{(i-1)(j+1)} + \gamma z_{(i+1)(j+1)}) \right. \\ &\left. + \sum_{\substack{l=(j+1) \\ l=1}}^{N_o} S_2 (\beta z_{il} + \delta z_{(i-1)(j+1)} + \delta z_{(i+1)(j+1)}) \right) + \hat{b} \end{aligned}$$

and an analogous form of $F_{i(j+1)}$. It is easy to see that $\sigma_j \mathbf{F}_i(\mathbf{Z}) = \mathbf{F}_i(\sigma_j \mathbf{Z})$ for all $j = 1 \dots N_o$, and for all $i = 1, \dots, N_a$; therefore the dynamics are equivariant under the action of S_{N_o} .

Elements $\rho \in D_{N_a}$ act on V by permuting the order of the agent vectors \mathbf{Z}_i in the total system vector $\mathbf{Z} = (\mathbf{Z}_1, \dots, \mathbf{Z}_{N_a})$. The generators of D_{N_a} are the reflection element ρ_1 which reverses the order of elements in \mathbf{Z} , and a rotation ρ_2 which cycles forward the vector by one element, mapping each element i to $i+1$ (and N_a to 1). Let $\mathbf{F}(\mathbf{Z}) = (\mathbf{F}_1(\mathbf{Z}), \dots, \mathbf{F}_{N_a}(\mathbf{Z}))$ and observe that $\rho_1 \mathbf{F}(\mathbf{Z}) = (\mathbf{F}_{N_a}(\mathbf{Z}), \mathbf{F}_{N_a-1}(\mathbf{Z}), \dots, \mathbf{F}_2(\mathbf{Z}), \mathbf{F}_1(\mathbf{Z}))$ and $\rho_2 \mathbf{F}(\mathbf{Z}) = (\mathbf{F}_{N_a}(\mathbf{Z}), \mathbf{F}_1(\mathbf{Z}), \mathbf{F}_2(\mathbf{Z}), \dots, \mathbf{F}_{N_a-1}(\mathbf{Z}))$. For compactness we leave out the full expression for $F_{ij}(\rho_p \mathbf{Z})$, and simply note that from writing it out we can conclude that $\rho_1 \mathbf{F}(\mathbf{Z}) = \mathbf{F}(\rho_1 \mathbf{Z})$ and $\rho_2 \mathbf{F}(\mathbf{Z}) = \mathbf{F}(\rho_2 \mathbf{Z})$, meaning the dynamics are equivariant under the action of D_{N_a} . \square

Propositions IV.1 and IV.2 reveal that symmetry in the opinion interaction topology constrains the number of parameters determining opinion network dynamics. In the maximally symmetric case, Proposition IV.1 shows that there are only four such parameters. The intra-agent opinion dynamics is entirely determined by the two weights α, β , while the inter-agent opinion dynamics is entirely determined by the two

weights γ, δ . When the agent symmetry group is smaller than S_{N_a} , e.g., D_{N_a} , the number of parameters determining the inter-agent opinion dynamics can increase but the sufficient condition in Proposition IV.1 shows there are again only two such parameters associated to any given inter-agent coupling graph with symmetry D_{N_a} , e.g., an undirected N_a -cycle. The same is true for different agent symmetry groups, or when some of the option symmetry is lost.

Model-independent analysis summarized in [33, Theorem 4.6 and Remark 4.7] predicts that in $(\Gamma_{N_o} \times \Gamma_{N_a})$ -equivariant opinion dynamics, with transitive Γ_{N_a} , opinionated steady-state solutions generically emerge from the neutral equilibrium $\mathbf{Z} = 0$ as bifurcations along the consensus space W_c or dissensus space W_d . Here, we illustrate how this model-independent analysis translates to predicting the behavior of the general model (5). We start by proving a general result providing the critical values of the control parameter for which bifurcations happens along the consensus or dissensus space for a generic (not necessarily symmetric) in-regular inter-agent coupling topology. We then leverage this result, Propositions IV.1 and IV.2, and the model-independent theory, to study the appearance of consensus and dissensus opinion formation in our general model.

Theorem IV.3 (Consensus and Dissensus). *Consider model (5) with $b_{ij} = \hat{b} \in \mathbb{R}$, $d_{ij} = d > 0$, $u_i = u \geq 0$, $A_{ii}^{jj} = \alpha \in \mathbb{R}$, $A_{ii}^{jl} = \beta \in \mathbb{R}$, $A_{ik}^{jj} = \gamma \tilde{a}_{ik}$, $A_{ik}^{jl} = \delta \tilde{a}_{ik}$, with $\gamma, \delta \in \mathbb{R}$ and $\tilde{a}_{ik} \in \{0, 1\}$ for all $i, k = 1, \dots, N_a$, $i \neq k$, and for all $j, l = 1, \dots, N_o$, $j \neq l$. Let $(\tilde{a}_{ik}) := \tilde{A} \in \mathbb{R}^{N_a \times N_a}$ and let J denote the Jacobian of this system evaluated at $\mathbf{Z} = 0$.*

A. Consensus. *Suppose the neutral equilibrium $\mathbf{Z} = \mathbf{0}$ is locally exponentially stable for $0 < u < u_c$ and $\ker J = W_c$ for $u = u_c$, i.e., a steady-state bifurcation occurs along the consensus space. If each node in the (directed or undirected) graph generated by \tilde{A} has in-degree K ,*

$$u_c = \frac{d}{\alpha - \beta + K(\gamma - \delta)} \quad (15)$$

and $(\alpha - \beta) > 0$, $(\gamma - \delta) > 0$. If \tilde{A} is a connected undirected graph, then more generally

$$u_c = \frac{d}{\alpha - \beta + \lambda_{max}(\gamma - \delta)} \quad (16)$$

where $\lambda_{max} > 0$ is the greatest eigenvalue of \tilde{A} and $(\alpha - \beta) > 0$, $(\gamma - \delta) > 0$.

B. Dissensus. *Suppose the neutral equilibrium $\mathbf{Z} = \mathbf{0}$ is locally exponentially stable for $0 < u < u_d$ and $\ker J = W_d$ for $u = u_d$, i.e., a steady-state bifurcation occurs along the dissensus space. If \tilde{A} is a connected undirected graph,*

$$u_d = \frac{d}{\alpha - \beta + \lambda_{min}(\gamma - \delta)} \quad (17)$$

where λ_{min} is the smallest eigenvalue of \tilde{A} and $(\alpha - \beta) > 0$, $(\gamma - \delta) < 0$.

Proof. Observe that $J = \mathcal{I} \otimes A_0 + \tilde{A} \otimes B_0$ where \otimes is the Kronecker product of matrices, $\mathcal{I} \in \mathbb{R}^{N_a \times N_a}$ is the identity matrix, and the matrices A_0, B_0 are defined as

$$A_0 = \begin{pmatrix} a & b & \dots & b \\ b & a & \dots & b \\ \vdots & \vdots & \ddots & \vdots \\ b & b & \dots & a \end{pmatrix}, \quad B_0 = \begin{pmatrix} c & e & \dots & e \\ e & c & \dots & e \\ \vdots & \vdots & \ddots & \vdots \\ e & e & \dots & c \end{pmatrix}$$

with $a = \frac{N_o - 1}{N_f}(-d + u\alpha - u\beta)$, $b = -\frac{1}{N_o - 1}a$, $c = \frac{N_o - 1}{N_o}u(\gamma - \delta)$, $e = -\frac{1}{N_o - 1}c$.

If $\ker J = W_c$ and all vertices of \tilde{A} have in-degree K , $(\mathcal{I} \otimes A_0 + \tilde{A} \otimes B_0)\mathbf{Z}_c = 0$ for any $\mathbf{Z}_c \in W_c$; using definition of W_c and some simple algebraic manipulation it follows that $(N_o - 1)(-d + u_c\alpha - u_c\beta + K u_c(\gamma - \delta)) = 0$ and (15) follows by solving for u_c .

The two matrices summed in the statement of J commute and we find that all eigenvalues of J correspond to $\mu_i = -d + u(\alpha - \beta) + u\lambda_i(\gamma - \delta)$ where λ_i is an eigenvalue of \tilde{A} . For a nondegenerate bifurcation to occur at some value of u it must be true that $\alpha - \beta > 0$ and $\lambda^*(\gamma - \delta) > 0$, with $\lambda^* = \lambda_i$ corresponding to the eigenvalue μ_i which crosses zero. The first eigenvalue to cross zero corresponds either to $\lambda^* = \lambda_{max}$ when $\gamma - \delta > 0$ or $\lambda^* = \lambda_{min}$ if $\gamma - \delta < 0$. From the properties of the eigenvectors of undirected connected graphs we deduce that if $\ker J = W_c$, $\lambda^* = \lambda_{max}$ and if $\ker J = W_c$, $\lambda^* = \lambda_{min}$, from which the final conclusions for A,B follow. \square

In general, it is difficult to compute the kernel of the Jacobian of model (5) and thus to apply Theorem IV.3. A basic result of equivariant bifurcation theory is that if the model has symmetries then, generically, $\ker J$ is an *irreducible representation* of the symmetry group [Vol 2 page...]. When the model symmetry group is $\Gamma = \Gamma_{N_a} \times \Gamma_{N_o}$ and Γ_{N_a} is transitive, then the only two irreducible representations of Γ on V are exactly the consensus and dissensus spaces (see [33, Remark 4.7]), which leads to the following corollary.

Corollary IV.3.1 (Parameter Conditions for Consensus and Dissensus). *Consider the $(S_{N_o} \times S_{N_a})$ -equivariant all-to-all network from Proposition IV.1 and the $(S_{N_o} \times D_{N_a})$ -equivariant undirected cycle network from Proposition IV.2. If $\alpha - \beta > 0$ and $\gamma - \delta > 0$, then, generically, opinion formation from the neutral equilibrium $\mathbf{Z} = 0$ occurs as a bifurcation along the consensus space for $u = u_c$, with $K = N_a - 1$ for the all-to-all network and $K = 2$ for the cycle. If $\alpha - \beta > 0$ and $\gamma - \delta < 0$, then, generically, opinion formation from the neutral equilibrium $\mathbf{Z} = 0$ occurs as a bifurcation along the dissensus space for $u = u_d$, with $\lambda_{min} - 1$ for the all-to-all network, $\lambda_{min} = -2$ for a cycle when N_a is even, and $\lambda_{min} = 2 \cos(\pi(N_a - 1)/N_a)$ when N_a is odd.*

Proof. Because both S_{N_a} and D_{N_a} are transitive, [33, Theorem 4.6 and Remark 4.7] ensure that generically $\ker J = W_c$ or $\ker J = W_d$. The result then follows directly from Theorem IV.3, noticing that, both in the S_{N_a} and D_{N_a} cases, the conditions $\alpha - \beta > 0$, $\gamma - \delta > 0$ ensures that $u_c > 0$ and either $u_d < 0$ (so dissensus bifurcations cannot happen for positive u) or $u_c < u_d$, so dissensus bifurcations happen after




Fig. 2: (a) Consensus in an all-to-all homogeneous network from Proposition IV.1, $\gamma = 0.2$, $\delta = -0.1$; (b) Dissensus $\gamma = -0.1$, $\delta = 0.2$; for $N_o = 2$, $N_a = 8$ (top) and $N_o = 3$, $N_a = 12$ (bottom). For all shown trajectories, the $\alpha = 0.2$, $\beta = 0.1$, $d = 1$, $u = 3$, $\hat{b} = 0$. Shown consensus and dissensus trajectories are initiated from the same random initial conditions. All parameters (including the communication weights $\alpha, \beta, \gamma, \delta$) were perturbed with small random additive perturbations drawn from a normal distribution with (a) variance 0.01, (b) variance 0.001. Opinion variables z_{ij} are mapped to the simplex for 3 options using the mapping in Corollary III.5.1 with $r = 1$ and $R = 2u$.

consensus bifurcations and thus play no opinion formation role. Conversely, for $\alpha - \beta > 0$, $\gamma - \delta < 0$, $u_d > 0$ and either $u_c < 0$ or $u_d < u_c$. \square

The parameter conditions in Theorem IV.3 and Corollary IV.3.1 can be read as follows: if agents cooperate ($\gamma > \delta$), opinions form along the consensus space; if agents compete ($\gamma < \delta$), opinions form along the dissensus space. The condition $(\alpha - \beta) > 0$ is satisfied when options are self-reinforcing ($A_{ii}^{jj} \geq 0$) and mutually exclusive ($A_{ii}^{jj} < 0$).

These highly symmetric cases provide the foundation to extend to the study and prediction of behavior in asymmetric cases much in the same way that the classical all-to-all and balanced cases do for linear and nonlinear consensus and synchronization dynamics [4], [47]–[49].

Remark IV.2 (Multistability). *An important consequence of Theorem IV.3 is the multistability of equilibria that arise when the system has symmetry. For example for the maximally symmetric system, whenever there exists a stable consensus equilibrium corresponding to the group favoring one of the options, due to the permutation symmetry between option there also exist $N_o - 1$ other stable equilibria corresponding to the group favoring each of the other options with the same magnitude of opinion. Analogously due to the symmetries of the group, stable dissensus equilibria appear in groups of $N_a N_o$ elements.*

Remark IV.3 (Robustness). *Consensus and dissensus solution branches predicted for the symmetric networks in Corollary IV.3.1 are a consequence of the Equivariant Branching Lemma [36, Section 1.4], and are made of hyperbolic equilibria. Their stability can be studied using the tools in [50, Section XIII.4] and [36, Section 2.3] although in high-dimensional cases the resulting computations might be intractable. Hyperbolicity of the equilibria gives rise to a notion of robustness of these solutions, due to general robustness of hyperbolic equilibria of dynamical systems. Introduction of small heterogeneous relative inputs and small asymmetries in the communication network will still result in a system with hyperbolic equilibria with the same stability properties near the consensus and dissensus spaces. Exact robustness bounds for asymptotically stable equilibria of (5) can be derived using similar methods to those used for general Hopfield networks in [51].*

An illustration of the formation of consensus and dissensus in an all-to-all network of agents evaluating 2 and 3 options, where small perturbations are added to all parameters including interactions weights, is provided in Fig. 2.

Remark IV.4 (Mode Interaction and Coexistence of Consensus and Dissensus). *When $\gamma = \delta$, there is mode interaction [37], and bifurcations along the consensus and dissensus spaces occur at the same critical value of u . The parameter regime is especially interesting because it allows for simultaneous stability of consensus and dissensus solutions, which can result in agents easily transitioning between consensus and dissensus in response to changing conditions. However, additional primary solution branches not captured by equivariant analysis can appear in this regime; classification of these for higher dimensional permutation symmetries remains an open problem.*

In the remainder of this section we show how the conditions for consensus and dissensus derived in Theorem IV.3 generalize, under certain conditions, to a network with N_c clusters.

Proposition IV.4 (Symmetric Clustered Dynamics). *Suppose in addition to satisfying assumptions of Theorem III.6, the parameters of (11) satisfy $u_p = u \geq 0$, $d_{pj} = d > 0$, $b_{pj} = \hat{b} \in \mathbb{R}$, and the interaction weights satisfy*

$$\alpha := \bar{B}_{pp}^{jj} + (N_p - 1)B_{pp}^{jj}, \quad \gamma := N_s B_{ps}^{jj}, \quad (18a)$$

$$\beta := \bar{B}_{pp}^{jl} + (N_p - 1)B_{pp}^{jl}, \quad \delta := N_s B_{ps}^{jl}, \quad (18b)$$

with $\alpha, \beta, \delta, \gamma \in \mathbb{R}$, for all $p, s = 1, \dots, N_c$, $j, l = 1, \dots, N_o$, $p \neq s$, $j \neq l$. Every trajectory of the dynamics (11) converges exponentially in time to a $N_c(N_o - 1)$ -dimensional attracting manifold \mathcal{E} defined by (9), the reduced dynamics on which are $(S_{N_o} \times S_{N_c})$ -equivariant. The associated clustered dissensus space along which solution branches emerge is

$$\hat{W}_d = \{(\mathbf{Z}_1, \dots, \mathbf{Z}_{N_a}) \mid \sum_{p=1}^{N_c} \frac{1}{N_p} \sum_{i \in \mathcal{I}_p} \mathbf{Z}_i = \mathbf{0}\}. \quad (19)$$

Proof. This follows directly from model reduction in Theorem III.6 and Proposition IV.1. \square




Fig. 3: Dynamics of a 3-cluster network with $N_1 = 2$, $N_2 = 3$, $N_3 = 4$. For all shown trajectories the weights in (18) satisfy $u = 4$, $d = 1$, $\hat{b} = 0$, $\bar{B}_{pp}^{jj} = B_{pp}^{jj}$, $\bar{B}_{pp}^{jl} = B_{pp}^{jl}$, $\alpha = 0.2$, $\beta = 0.1$. (a) Consensus, $\gamma = 0.2$, $\delta = -0.1$; (b) Dissensus $\gamma = -0.1$, $\delta = 0.2$. Small random perturbations drawn from a normal distribution with variance 0.01 were added to all of the model parameters (including the communication weights $\alpha, \beta, \gamma, \delta$). Opinion variables z_{ij} are mapped to the simplex for 3 options using the mapping in Corollary III.5.1 with $r = 1$ and $R = 2u$.

An illustration of the formation of consensus and dissensus in a three cluster network evaluating 3 options, where small perturbations are added to all parameters including interactions weights, is provided in Fig. 3.

V. SPECIALIZATION TO MODELS IN THE LITERATURE

In this section we show how the general model (5) for $N_o = 2$ options specializes to well-studied nonlinear and linear consensus models in the literature. We also show that network conditions, e.g., structural balance, proved to be necessary for clustering in linear models, e.g., for bi-partite consensus [5], are not necessary when there is a saturating nonlinearity.

For $N_o = 2$, the opinion state of agent i is one-dimensional: $\mathbf{Z}_i = (z_{i1}, z_{i2})$, with $z_{i1} + z_{i2} = 0$. We define $x_i = z_{i1} = -z_{i2}$ as agent i 's opinion. Then, opinion dynamics (5) reduce to

$$\begin{aligned} \dot{x}_i = & -d_i x_i + \frac{1}{2} u_i \left(S_1 \left(\sum_{k=1}^{N_a} A_{ik}^{11} x_k \right) - S_1 \left(- \sum_{k=1}^{N_a} A_{ik}^{22} x_k \right) \right) \\ & + S_2 \left(- \sum_{k=1}^{N_a} A_{ik}^{12} x_k \right) - S_2 \left(\sum_{k=1}^{N_a} A_{ik}^{21} x_k \right) + b_i^\perp \end{aligned} \quad (20)$$

where $b_i := b_{i1} = -b_{i2}$, and $d_i = \frac{1}{2}(d_{i1} + d_{i2})$. Let the network opinion state be $\mathbf{x} = (x_1, \dots, x_{N_a}) \in \mathbb{R}^{N_a}$ and vector of relative inputs be $\mathbf{b}^\perp = (b_1^\perp, \dots, b_{N_a}^\perp) \in \mathbb{R}^{N_a}$.

We first show how (20) reduces to the nonlinear model of consensus studied in [15]–[17].

Theorem V.1. *For N_a agents and $N_o = 2$ options, let $A_{ii}^{jj} = A_{ii}^{jl} = A_{ik}^{jl} = 0$, $A_{ik}^{jj} = \gamma_{ik} \in \mathbb{R}$, $u_i = u \geq 0$, and*

$$d_{ij} = \sum_{k=1}^{N_a} \sum_{l=1}^{N_o} |A_{ik}^{jl}| = \sum_{k=1}^{N_a} |\gamma_{ik}| := d_i, \quad (21)$$

for $i, k = 1, \dots, N_a$, $i \neq k$, $j, l = 1, \dots, N_o$, $j \neq l$. Then, opinion dynamics (5) reduce to

$$\dot{x}_i = -d_i x_i + u \hat{S}_1 \left(\sum_{k=1}^{N_a} \gamma_{ik} x_k \right) + b_i^\perp, \quad (22)$$




Fig. 4: For small initial conditions, the Altafini model (23) (left) approximates the response of the nonlinear model (22) (right) with $N_a = 3$, $u = 1$, $b_i^\perp = 0$, and $\hat{S}_1 = \tanh$. As expected, clustered behavior is observed for a structurally balanced graph for both models. Simulations use the structurally balanced adjacency matrix from [5, Example 1]. The same small random initial conditions are used for both models.

where $i \neq k$ and $\hat{S}_p(x) = \frac{1}{2}(S_p(x) - S_p(-x))$, $p \in \{1, 2\}$, are odd sigmoids and where $x_i := z_{i1} = -z_{i2}$, $b_i := b_{i1} = -b_{i2}$, and $d_i = \frac{1}{2}(d_{i1} + d_{i2})$. In particular, (5) specializes to the model of nonlinear consensus dynamics studied in [15]–[17].

Proof. Since $N_o = 2$, (5) reduced to (20). By the assumptions, (20) further reduces to (22). For $\gamma_{ik} \geq 0$, dynamics (22) are equivalent to the nonlinear model [15]–[17]. \square

We next show how (20) reduces to the ‘‘Altafini’’ model of linear consensus with antagonistic interconnections [5].

Proposition V.2. *The Altafini model [5] is recovered from (5) as a linearization of (22) about $\mathbf{x} = 0$ with $u = 1$ and $b_i^\perp = 0$:*

$$\dot{\mathbf{x}} = -\mathcal{L}\mathbf{x}. \quad (23)$$

$\mathcal{L} = \mathcal{D} - \mathcal{A}$ is the signed graph Laplacian matrix, with degree matrix $\mathcal{D} = \text{diag}(d_i) \in \mathbb{R}^{N_a \times N_a}$ and adjacency matrix $\mathcal{A} \in \mathbb{R}^{N_a \times N_a}$ having entries that are the inter-agent weights γ_{ik} .

Proof. By Theorem V.1, (5) reduces to (22). It is shown in [16] that the linearization of (22) recovers the Altafini model. \square

In the Altafini model, a clustered disagreement outcome referred to as *bipartite consensus* arises for the network only if the communication graph among agents is structurally balanced (i.e. \mathcal{L} has a zero eigenvalue); see [5, Theorem 2]. Multi-option generalizations of linear bipartite consensus protocols also require structural balance for clustered behavior [30]. By Proposition V.2, this property is recovered locally in the nonlinear model (22), as illustrated in Fig. 4. However, in the light of Theorems III.6 and IV.4, we have the following corollary, illustrated in Fig. 5.

Corollary V.2.1. *A structurally balanced network is not necessary for opinion clustering in the nonlinear model. 5.*

VI. DYNAMIC FEEDBACK AND TUNABLE SENSITIVITY

We have established that existence of consensus and dissensus equilibria and multistability of opinion formation outcomes arise from bifurcations of the general opinion dynamic




Fig. 5: Altafini model (23) (left) and nonlinear model (22) with $N_a = 5$, $u = 5$, $b_i^\perp = 0$, $\hat{S}_1 = \tanh$, same random initial conditions, and same adjacency matrix given by $\gamma_{ik} = -1$ for $i, k \in \mathcal{I}_p$, $i \neq k$, and $\gamma_{ik} = -2$ for $i \in \mathcal{I}_p$, $k \in \mathcal{I}_s$, $p \neq s$ for clusters with indices $\mathcal{I}_1 = \{1, 2\}$ and $\mathcal{I}_2 = \{3, 4, 5\}$. Because the adjacency matrix is not structurally balanced, the Altafini model converges to the neutral solution. The nonlinear model, however, converges to a stable dissensus state, as predicted. The parameters satisfy sufficient conditions of Theorem III.6 for trajectories to converge to a 2-dimensional manifold. On this manifold, the clustered dynamics are $(S_2 \times S_2)$ -equivariant with interaction weights $\alpha = 0$, $\beta = -2$, $\gamma = 0$, $\delta = -2$, which satisfy the conditions for dissensus from Theorem IV.3.

model (5). In this section we explore how *ultra-sensitivity to inputs* b_{ij} , *robustness to changes in inputs*, *cascade dynamics*, and *transitions between consensus and dissensus solutions all arise as a consequence of the multistability*. With the addition of dynamic state feedback for model parameters in (5), the opinion formation process can reliably amplify arbitrarily small relative inputs b_{ij}^\perp , reject small changes in input as unwanted disturbance, facilitate an opinion cascade even if only one agent gets an input, and enable groups to move easily between consensus and dissensus. Moreover, the choice of feedback design parameters determine implicit thresholds that make all of these behaviors are tunable.

In Section VI-A we propose a dynamic state feedback law for the attention parameter u_i in the general model (5). We then focus on $N_o = 2$ and show how the dynamic feedback law enables tunable sensitivity and robustness (Section VI-B), tunable cascade dynamics (VI-C), and tunable consensus/dissensus transitions (VI-D). To further simplify the exposition, we specialize the analysis to the case in which the intra- and inter-agent coupling weights do not depend on the option, i.e., we let $A_{ii}^{jj} := \alpha_i$, $A_{ii}^{ll} := \beta_i$, $j \neq l$, $A_{ik}^{jj} := \gamma_{ik}$, $A_{ik}^{ll} := \delta_{ik}$, for $j, l = 1, 2$, $j \neq l$, $i, k = 1, \dots, N_a$, $i \neq k$ such that dynamics (20) become

$$\dot{x}_i = -d_i x_i + b_i^\perp + u_i \left(\hat{S}_1 \left(\alpha_i x_i + \sum_{k=1}^{N_a} \gamma_{ik} x_k \right) - \hat{S}_2 \left(\beta_i x_i + \sum_{k=1}^{N_a} \delta_{ik} x_k \right) \right) \quad (24)$$

where $x_i, d_i, b_i^\perp, \hat{S}_p$ are defined as in Section V.

A. Dynamic State Feedback Law for Attention

In design of the feedback control laws we take inspiration from neuronal gating mechanisms in biological and artificial neural networks. In these systems, modulation of gain parameters is central to regulating real-time information processing [52] and implementing optimal decision-making strategies [42], [53], [54]. Consensus cascades in a neural network opinion model using a similar state feedback mechanism were recently explored in [55] for two options.

We augment the opinion dynamics model (5) by introducing feedback dynamics on the attention parameter u_i for each agent i , in the form of a saturated nonlinear leaky integrator:

$$\begin{aligned} \tau_u \dot{u}_i &= -u_i + S_u \left(\frac{1}{N_o^2} \sum_{j=1}^{N_o} \sum_{k=1}^{N_a} \sum_{l=1}^{N_o} \left(\bar{A}_{ik}^{jl} z_{kl} \right)^2 \right) \\ &= -u_i + S_u \left(\frac{1}{N_o^2} \|\bar{A}_i \mathbf{Z}\|^2 \right), \end{aligned} \quad (25)$$

where $\bar{A}_i \in \mathbb{R}^{N_o \times N_a N_o}$ with \bar{A}_{ik}^{jl} the entry in row j , column $N_a(k-1) + l$. Here, $\tau_u > 0$ is the time scale. S_u is a smooth saturating function, satisfying $S_u(0) = 0$, $S_u(y) \rightarrow u_{sat} > 0$ as $y \rightarrow \infty$, $S'_u(y) > 0$ for all $y \in \mathbb{R}$, and $S''_u(y) < 0$ for all $y > y_m$, and $y_m > 0$. We can always decompose S_u as

$$S_u(y) = u_f \left(F(g(y - y_m)) - F(-gy_m) \right), \quad (26)$$

$F: \mathbb{R} \rightarrow [0, 1]$ a smooth monotone sigmoid with $F'(x) > 0$ for all $x \in \mathbb{R}$, $F'(x) = F'(-x)$, $F''(x) < 0$ for $x > 0$. The parameter g defines the slope of S_u . We will show how u_f and y_m are sensitivity and robustness tuning design parameters:

- design parameter u_f defines the upper bound of S_u ;
- design parameter y_m defines the midpoint of S_u .

The input to S_u in (25) is proportional to the magnitude of the network opinion state that influences the opinion of agent i . The weights \bar{A}_{ik}^{jl} can be real or binary $\{0, 1\}$. \bar{A}_{ik}^{jl} can be related to the corresponding interaction weights A_{ik}^{jl} in (5), or independently defined. The latter case allows agents to exert influence by exciting their neighbors to form an opinion without influencing what opinion they form.

For all simulations we define S_u by (26) with $F(x) = 1/(1 + e^{-x})$. Large inputs saturate to $u_{sat} = u_f e^{gy_m}/(1 + e^{gy_m})$. For $gy_m > 4$, u_f is a good approximation of u_{sat} .

Remark VI.1 (Well-Definedness of Opinion Dynamics with Feedback). *Observe that $\dot{u}_i(0) < 0$ for $u_i(0) \geq u_{sat}$ and $\dot{u}_i(0) > 0$ for $u_i(0) \leq 0$. It follows that the set $\{0 \leq u_i \leq u_{sat}\}$ is attractive and forward invariant, and $|\dot{u}_i|$ is uniformly bounded in $\{0 \leq u_i \leq u_{sat}\}$. Thus, boundedness of $\mathbf{Z}(t)$ in the presence of attention feedback dynamics follows along the same lines as the proof of Theorem III.5. So, the attention feedback dynamics do not modify invariance of the simplex product \mathcal{V} and the coupled dynamics (5),(25) are well-defined.*

B. Tunable Sensitivity and Robustness

We first examine the opinion dynamics (24) for uncoupled agents, i.e. $\gamma_{ik} = \delta_{ik} = 0$, which are

$$\dot{x}_i = -d_i x_i + u_i \hat{S}(x_i) + b_i^\perp := q(x_i, u_i, b_i^\perp) \quad (27)$$

with $\hat{S}(x_i) := \hat{S}_1(\alpha_i x_i) - \hat{S}_2(\beta_i x_i)$. The following assumption ensures that \hat{S} is a monotonically increasing sigmoid.

Assumption 1. $\alpha_i > 0 > \beta_i$ for all $i = 1, \dots, N_a$.

Since Assumption 1 is satisfied for self-reinforcing, mutually exclusive options, we let it hold for the rest of the paper. In all simulations we let $\hat{S}_1(x) = \tanh(x)$ and $\hat{S}_2(x) = \frac{1}{2} \tanh(2x)$. The following proposition classifies the possible steady-state behaviors of (27); see Fig. 6 for an illustration.

Proposition VI.1 (Opinion Dynamics of Uncoupled Agents). *For the one-dimensional dynamics (27) of the opinion state of agent i , the following hold:*

A. *For $b_i^\perp = 0$ and $0 \leq u_i \leq u_i^*$ with $u_i^* = \frac{d_i}{\alpha_i - \beta_i}$ the unopinionated state $x_i = 0$ is globally asymptotically stable and locally exponentially stable for $0 < u_i < u_i^*$;*

B. *For $b_i^\perp = 0$, the bifurcation problem $g(x_i, u_i, 0) = 0$ has a pitchfork singularity at $(x_i^*, u_i^*) = \left(0, \frac{d_i}{\alpha_i - \beta_i}\right)$. For $u_i > u_i^*$ the neutral equilibrium $x_i = 0$ loses stability and two additional branches of locally exponentially stable equilibria $x_i = \pm x_i^s$ emerge, implicitly defined by $0 = -d_i x_i^s + u_i \hat{S}(x_i^s)$;*

C. *For $b_i^\perp \neq 0$, $q(x_i, u_i, b_i^\perp)$ is a 1-parameter unfolding of the symmetric pitchfork. For u_i in a small neighborhood of u_i^* , a single locally exponentially stable equilibrium x_i^s exists that has the same sign as b_i^\perp . For sufficiently large values of u_i away from the singularity u_i^* , a second locally exponentially stable equilibrium exists which has the opposite sign as b_i^\perp .*

Proof. Uncoupled agent dynamics (27) are equivalent to a one-dimensional instance of the nonlinear consensus model studied in [16]. Conclusions A – C follow from [16, Theorem 1]. \square

We next establish how the shape of the unfolded bifurcation diagram (see Fig. 6, left) described in Proposition VI.1-C changes with input b_i^\perp (see Fig. 6, right).

Lemma VI.2 (Input Response Without Feedback). *Let x_i^s be a hyperbolic equilibrium of (27) with input b_i^\perp and $u_i \geq 0$. Then $\frac{\partial x_i^s}{\partial b_i^\perp} > 0$ if x_i^s is stable, and $\frac{\partial x_i^s}{\partial b_i^\perp} < 0$ if it is unstable.*

Proof. Implicit differentiation of $q(x_i^s, u_i, b_i^\perp) = 0$ by b_i^\perp at a constant u_i yields

$$\frac{\partial x_i^s}{\partial b_i^\perp} = \frac{1}{d_i - u_i \hat{S}'(x_i^s)} = -\frac{1}{\lambda} \quad (28)$$

where $\lambda \in \mathbb{R}$ is the eigenvalue of the linearization of (27) evaluated at an equilibrium. At any stable equilibrium of (27), $\lambda < 0$ and therefore $\frac{\partial x_i^s}{\partial b_i^\perp} > 0$. The unstable case follows. \square

We now introduce the attention feedback dynamics for u_i (25). Let the feedback weights be

$$\bar{A}_{ik}^{11} = \bar{A}_{ik}^{12} = \bar{A}_{ik}^{21} = \bar{A}_{ik}^{22} := \bar{a}_{ik} \in \{0, 1\} \quad (29)$$

with $\bar{a}_{ik} = 1$ whenever u_i is influenced by the state of agent k . The attention feedback law (25) then simplifies to

$$\tau_u \dot{u}_i = -u_i + S_u \left(\sum_{k=1}^{N_a} (\bar{a}_{ik} x_k)^2 \right). \quad (30)$$




Fig. 6: Bifurcation diagrams of (27) with $\alpha_i = 2, \beta_i = -1, d_i = 1$ with no bias (left) and a positive bias (right). Black lines plot the steady-state solutions (nullclines) and gray arrows are streamlines showing direction of the flow.

When agent i is completely decoupled from all neighbors, i.e., $\bar{a}_{ii} = 1$ and $\bar{a}_{ik} = 0$ for all $k \neq i$, (30) reduces to

$$\tau_u \dot{u}_i = -u_i + S_u(x_i^2) := h(x_i, u_i). \quad (31)$$

As illustrated in Fig. 7, the equilibria of system (27),(31) can be visualized as intersections of the x_i -nullcline $q(x_i, u_i, b_i^\perp) = 0$ (black solid) and u_i -nullcline $h(x_i, u_i) = 0$ (red dashed). Proposition VI.1 defines the shape of the x_i -nullcline; see Fig. 6 (right); Lemma VI.2 describes how the shape changes with input b_i^\perp . Let agent i be *strongly opinionated* when it is opinionated and its attention is close to its saturation value, i.e., $u_i \simeq u_{sat}$.

Tunable sensitivity of opinion formation to input b_i^\perp can then be understood by comparing the plots of Fig. 7, where the trajectory for agent i is plotted on the left for $b_i^\perp = 0.5$ and on the right for $b_i^\perp = 1.5$. For the given parameters and $b_i^\perp = 0.5$, the nullclines intersect at three points. Starting at the origin, the opinion state is attracted to the point corresponding to an unopinionated equilibrium: agent i *rejects the input* $b_i^\perp = 0.5$ and does not form a strong opinion. For the same parameters and $b_i^\perp = 1.5$, the nullclines intersect at only one point, corresponding to a strongly opinionated equilibrium. Thus, for the same initial condition, agent i *accepts the input* $b_i^\perp = 1.5$ and forms a strong opinion. The implicit *sensitivity threshold* that distinguishes rejected from accepted input magnitude can be tuned by design parameter y_m .

Tunable robustness of opinion formation to changes in input b_i^\perp can be understood by comparing the sequence of plots in Fig. 8(a) to the sequence of plots in Fig. 8(b). In (a) and (b) the plot on the left shows agent i forming a strong opinion in the direction of the input $b_i^\perp = 1$. In (a) and (b) the plot on the right shows what happens to agent i 's opinion when the input changes to $b_i^\perp = -1$, i.e., an input that is in the opposite direction of the original input. In (a), when $u_f = 1$, agent i *accepts the change of input* and forms a strong opinion in the direction of the new input. In (b), when $u_f = 2.5$, agent i *rejects the change of input* and retains a strong opinion in the direction of the original input. The implicit *robustness threshold* that distinguishes rejected from accepted changes in input can be tuned by design parameter u_f .

To develop systematic tools for tuning with design parameters u_f and y_m , we first develop some geometric intuition




Fig. 7: Sensitivity of opinion formation to input magnitude. Trajectories of (27), (31) with $g = 10, y_m = 1, u_f = 2, \alpha_i = 2, \beta_i = -1, d_i = 1, \tau_u = 1$ for $b_i^\perp = 0.5$ (left) and $b_i^\perp = 1.5$ (right). Initial state $(u_i(0), x_i(0)) = (0, 0)$ is a blue circle, and final state is a yellow diamond. Nullclines of (27) are black solid lines. Nullclines of (31) are red dashed lines. Gray arrows show streamlines of the flow. Color scale is time.




Fig. 8: Robustness of opinion formation to changes in input. Trajectories of (27), (31) with $g = 10, y_m = 1, \alpha_i = 2, \beta_i = -1, d_i = 1, \tau_u = 1$. (Left) Input is $b_i^\perp = 1$, initial state $(u_i(0), x_i(0)) = (0, 0)$ is a blue circle, and final state is a cyan diamond. (Right) Input changes to $b_i^\perp = -1$, initial state is final state on left and final state is yellow square. (a) $u_f = 1$, and agent changes opinion in direction of new input. (b) $u_f = 2.5$, and agent retains opinion in original direction. Nullclines, streamlines, and time are drawn as in Fig. 7.

about trajectories of the coupled opinion and attention dynamics (27),(31).

Lemma VI.3 (Trapping Region). *The quadrant $u_i \geq 0, x_i \geq 0$ is forward-invariant for the coupled dynamics (27),(31) whenever $b_i^\perp \geq 0$. Similarly, the quadrant $u_i \geq 0, x_i \leq 0$ is forward-invariant whenever $b_i^\perp \leq 0$.*

Proof. This follows from observation that $\dot{u}_i \geq 0$ whenever $u_i = 0$ and $\dot{x}_i = b_i^\perp$ whenever $x_i = 0$. \square

Lemma VI.4 (Flow Monotonicity). *The flow generated by vector field (27),(31), restricted to the forward-invariant quad-*

rants from Proposition VI.3, is monotone in the sense of [56].

Proof. Consider $b_i^\perp > 0$ and restrict the system to the positive quadrant $u_i \geq 0, x_i \geq 0$. In this quadrant $\frac{\partial q}{\partial u_i} \geq 0$ and $\frac{\partial h}{\partial x_i} \geq 0$, and the proposition follows by [56, Chapter 3 Proposition 1.1, Remark 1.1]. The result for $b_i^\perp < 0$ and the negative quadrant $u_i \geq 0, x_i \leq 0$ follows analogously. \square

Lemmas VI.3, VI.4 show how agent i 's input picks out the trapping region for its dynamics. The trajectories of attention and opinion in the negative quadrant, $u_i \geq 0, x_i \leq 0$ for $b_i^\perp < 0$, are mirror-symmetric to trajectories in the positive quadrant $u_i \geq 0, x_i \geq 0$ for $b_i^\perp > 0$. Without loss of generality, we can thus assume $b_i^\perp \geq 0$ in the statement of the main results and in the geometric illustrations.

The equilibrium opinions x_i^s are solutions of

$$0 = -d_i x_i^s + S_u((x_i^s)^2) \hat{S}(x_i^s) + b_i^\perp, \quad (32)$$

and, in the forward-invariant quadrant determined by the sign of b_i^\perp , they all belong to the monotone, continuous branch of the x_i -nullcline (Fig. 6). This branch is made of hyperbolic equilibria of (27), and thus Lemma VI.2 applies. It follows that if slope g in S_u is sufficiently large, then, generically, the x_i and u_i -nullclines intersect three times (when $|b_i^\perp|$ is small, Fig. 7 left) or one time (when $|b_i^\perp|$ is large, Fig. 7 right) in the forward-invariant quadrant.

Fig. 9 plots x_i^s for four different parameter combinations. For sufficiently large g , (32) exhibits two fold singularities for $x_i^s \geq 0$ as b_i^\perp is varied. The right fold occurs for positive values of b_i^\perp and corresponds to the annihilation of the unopinionated equilibrium with a saddle point. The left fold occurs for negative values of b_i^\perp and corresponds to the annihilation of the strongly opinionated equilibrium with a saddle point. Equilibria and fold singularities for $x_i^s \leq 0$ are obtained by reflecting Fig. 9 along the $x_i^s = 0$ axis. We formalize this geometric discussion in Assumption 2, which we let hold for the rest of the paper.

Assumption 2. Design parameter g in S_u of (26) is sufficiently large so that (32) has two fold points in each half-plane of the (b_i^\perp, x_i^s) plane. Let $(\pm b_{LF}, \pm x_{LF})$ be the coordinates of left folds and $(\pm b_{RF}, \pm x_{RF})$ be the coordinates of the right folds, where $x_{RF} < \sqrt{y_m} < x_{LF}$, $b_{LF} < b_{RF}$, $0 < b_{RF}$.

Fold points of the opinion plus attention dynamics are important because they determine the appearance and disappearance of unopinionated and strongly opinionated equilibria. The fact that the fold locations depend on the model parameters as illustrated in Fig. 9 is proved in the following theorem.

Theorem VI.5 (Tunable Fold Point Location). *Consider the system (27),(31). Suppose $u_{sat} > u_i^* = \frac{d_i}{\alpha_i - \beta_i}$. Then, at the two fold points of (32), $\frac{\partial b_p}{\partial y_m} > 0$ and $\frac{\partial b_p}{\partial u_f} < 0$, $p \in \{RF, LF\}$. Further, the distance between the two folds increases with u_f : $\frac{\partial}{\partial u_f}(b_{LF} - b_{RF}) > 0$.*

Proof. Let the right hand side of (32) be $f_1(x_i^s, b_i^\perp)$. The expressions $f_1(x_p, b_p) = 0$ and $\frac{\partial f_1}{\partial x_p} := f_2(x_p, b_p) = 0$ for $p \in \{RF, LF\}$ together define the two fold points. The fold




Fig. 9: Solutions of (32). $g = 10$, $d_i = 1$, $\alpha_i = 2$, $\beta_i = -1$. (a) $u_f = 2.5$, $y_m = 1$ (solid), $y_m = 3$ (dashed). (b) $y_m = 2.5$, $u_f = 1$ (solid), $u_f = 2.5$ (dashed).

equations define x_p and b_p as implicit functions of y_m . Using the implicit function theorem, we find $\frac{\partial b_p}{\partial y_m} = -\frac{\partial f_1}{\partial y_m}$ at a fold:

$$\frac{\partial b_p}{\partial y_m} = u_f g \left(F'(g((x_p)^2 - y_m)) - F'(-gy_m) \right) \hat{S}(x_p) > 0.$$

Analogously, the fold equations define x_p and b_p as implicit functions of u_f . Using the implicit function theorem and the monotonicity of F , we find $\frac{\partial b_p}{\partial u_f} = -\frac{\partial f_1}{\partial u_f}$ at a fold:

$$\frac{\partial b_p}{\partial u_f} = - \left(F(g((x_p)^2 - y_m)) - F(-gy_m) \right) \hat{S}(x_p) < 0.$$

That $\frac{\partial b_{LF}}{\partial u_f} - \frac{\partial b_{RF}}{\partial u_f} > 0$ follows from the above expression and monotonicity of F and \hat{S} since $x_{RF} < x_{LF}$. \square

The following theorem shows how uncoupled agents with attention feedback dynamics can become strongly opinionated in response to a sufficiently strong input b_i^\perp . It also shows that the sensitivity threshold, which distinguishes between small and large inputs, can be tuned by tuning the right fold location according to Theorem VI.5.

Theorem VI.6 (Input Response for Uncoupled Agents with Attention Dynamics). *Let $u_{sat} > u_i^* = \frac{d_i}{\alpha_i - \beta_i}$. There exists $\epsilon > 0$ such that if $\|(x(0), u(0))\| < \epsilon$, the following hold for opinion and attention dynamics (27),(31):*

- For all $b_i^\perp \neq 0$, $(x_i(t), u_i(t))$ converges asymptotically to a nonzero equilibrium (x_i^s, u_i^s) with $u_i^s > 0$ and x_i^s of the same sign as b_i^\perp . The value of x_i^s is the value of the continuous solution branch in the pitchfork unfolding of (27) at $u_i = u_i^s$;
- There exist constants $0 < b_0 < b_{RF}$ and $\gamma_0 \geq 0$ such that for all $|b_i^\perp| \leq b_0$, there exists an equilibrium (x_i^s, u_i^s) with $\|(x_i^s, u_i^s)\| \leq \gamma_0 |b_i^\perp|$, and $\|(x_i(t) - x_i^s, u_i(t) - u_i^s)\| \leq Ae^{-t/\tau}$ for some $A, \tau > 0$.
- For any $\nu > 0$, there exists a threshold $b_{th} \geq b_{RF} > 0$ such that if $b_i^\perp > b_{th}$, then there exists a unique, locally exponentially stable equilibrium (x_i^s, u_i^s) in the quadrant $x_i, u_i \geq 0$, with $|u_i^s - u_{sat}| < \nu$. Moreover, $\lim_{t \rightarrow \infty} \|(x_i(t) - x_i^s, u_i(t) - u_i^s)\| = 0$.

Proof. A. By Lemma VI.4, the flow is monotonic. All trajectories are bounded, and convergence to an equilibrium in the positive quadrant follows from [56, Chapter 3 Theorem

2.2]. By Proposition VI.1-C, the equilibrium values x_i^s on the continuous branch of solutions are of the same sign as b_i^\perp .

B. When $b_i^\perp = 0$, the Jacobian of (27),(31) at equilibrium $(x_i^s, u_i^s) = (0, 0)$ has eigenvalues $-d_i, -1$. Thus, $(0, 0)$ is exponentially stable. Both $\frac{\partial q}{\partial b_i^\perp}$ and $\frac{\partial h}{\partial b_i^\perp}$ are bounded uniformly in time. The conclusion follows by [46, Corollary 5.1], definition of small-signal finite-gain stability [46, Definition 5.2], and monotonicity of the flow.

C. After the right fold, i.e., for $b_i^\perp > b_{RF}$ the only equilibrium in the quadrant $x_i, u_i \geq 0$ is the strongly opinionated equilibrium. By Lemma VI.2, the x_i -coordinate of this equilibrium can be made arbitrarily large by increasing b_i^\perp and thus $u_i^s = S_u((x_i^s)^2)$ can be made arbitrarily close to its saturation value u_{sat} . \square

Remark VI.2 (Sensitivity Threshold of Opinion Formation). *The coordinate of the right fold b_{RF} in (32) (Fig. 9) corresponds to the minimum threshold b_{th} from Theorem VI.6. Thus, b_{RF} indicates how sensitive the agent is to input: agent i will strongly amplify input $b_i^\perp > b_{RF}$ no matter how small, and will only slightly amplify $b_i^\perp \leq b_{RF}$; see Fig. 7.*

Remark VI.3 (Robustness Threshold of Opinion Formation). *The coordinate of the left fold b_{LF} in (32) (Fig. 9) indicates how robust the opinion formation is to changes in input. Suppose agent i becomes strongly opinionated with input $b_i^\perp = b_1 > 0$, and then the input switches to $b_i^\perp = b_2 < b_1$. Due to the hysteresis shown in Fig. 9, as long as $b_2 > b_{LF}$, agent i will remain strongly opinionated in the direction of b_1 , even if the b_2 is in favor of the alternative option; see Fig. 8.*

Theorems VI.5 and VI.6 show how design parameters u_f, y_m have complementary roles in tuning the system response: y_m tunes the sensitivity threshold and u_f tunes the robustness threshold. Increasing y_m shifts b_{RF} to the right, increasing the sensitivity threshold above which inputs will affect the opinion formation process. Increasing u_f has the primary effect of shifting b_{LF} to the left, thereby rendering the opinion formation more robust to input fluctuations. See Figs. 8 and 9 for illustration.

C. Tunable Cascade Dynamics

We next examine the opinion dynamics (27) for uncoupled agents, and introduce agent coupling in the attention feedback dynamics (30). Before showing how this leads to tunable cascade dynamics, we infer properties of opinion formation of a homogeneous group with a static u_i .

Assumption 3 (Homogeneous Agents). *The system parameters in a group of N_a agents with opinion dynamics (27) satisfy $\alpha_i = \alpha > 0$, $\beta_i = \beta < 0$, $d_i = d > 0$ for all $i = 1, \dots, N_a$.*

Proposition VI.7 (Properties of Homogeneous Uncoupled Agents). *The following hold for N_a agents with opinion dynamics (27), Assumption 3 and $u_i = u$:*

A. *When $b_i^\perp = 0$ for all $i = 1, \dots, N_a$, $\mathbf{x} = \mathbf{0}$ is a globally asymptotically stable equilibrium $0 \leq u \leq u^* := \frac{d}{\alpha - \beta}$ and locally exponentially stable for $0 \leq u < u^*$.*

B. *When $b_i^\perp = 0$ for all $i = 1, \dots, N_a$, and for $u > u^*$, the system has 2^{N_a} exponentially stable equilibria.*

C. *When $b_i^\perp \neq 0$ for exactly N_b of the N_a agents, the system has $2^{(N_a - N_b)}$ exponentially stable equilibria when $u^* < u < u_1$, and 2^{N_a} exponentially stable equilibria when $u > u_2$ for some $u_1 < u_2$.*

Proof. A, B, and C directly follow from corresponding parts A,B,C in Proposition VI.1 and combinatorics. \square

Each nonzero stable equilibrium in Proposition VI.7 corresponds to a unique allocation of agents into two subgroups, each subgroup favoring one of the options. Agents in different subgroups have opposite-sign opinion states x_i . When the attention parameter u is above but still close in value to its threshold u^* , the group opinion is sensitive to the inputs: agents with input always favor the option of the same sign and b_i^\perp agents without input favor either of the two.

Remark VI.4 (Mode Interaction and Control). *The completely decoupled system $\gamma = \delta = 0$ is a special instance of the mode interaction regime $\gamma = \delta$ of the opinion dynamics, as discussed in Remark IV.4. Proposition VI.7 shows how this parameter regime may be particularly useful for control applications. In this regime it is possible for stable consensus and dissensus solutions to coexist, and the additional branches of stable equilibria that may arise can provide a richer catalogue of behaviors available for system design. For example, the additional stable equilibria of Proposition VI.7 describe the different ways a group of uncoupled homogeneous N_a agents can be distributed across two options. Availability of such versatile and interpretable configurations of opinions can prove useful for applications involving dynamic task allocation.*

Now we consider a network of agents with coupling in the attention dynamics defined by (30). We define the *region of influence of agent i* in the network as the set of all nodes k to which there exists a directed path from node i in the graph with adjacency matrix \bar{A}_{ik} defined by $\bar{a}_{ik} \in \{0, 1\}$ as in (29). In the following theorem we show that it is enough for a single agent with sufficient influence in the network to receive a strong input in order for the entire network of agents to become strongly opinionated through a cascade.

Theorem VI.8 (Coupled Attention Dynamics and Opinion Cascades). *Consider N_a agents with opinion and attention dynamics governed by (27),(30), attention dynamics coupling defined by \bar{A}_{ik} as in (29). Let Assumption 3 hold. Assume $b_i^\perp \neq 0$ for all $i = 1, \dots, N_a$. Then, there exists $\epsilon > 0$ such that if $\|(\mathbf{x}(0), \mathbf{u}(0))\| < \epsilon$, the following hold:*

A. *The system asymptotically converges to a nonzero equilibrium point $(\mathbf{x}^s, \mathbf{u}^s)$ with $u_i^s > 0$ and each x_i^s of the same sign as the input b_i^\perp for all $i = 1, \dots, N_a$. The value of x_i^s is the value of the continuous solution branch in the pitchfork unfolding of (27) at $u_i = u_i^s$;*

B. *There exist constants $b_0 \in \mathbb{R}^{>0}$ and $\gamma_0 \in \mathbb{R}^{\geq 0}$ such that for all $\|\mathbf{b}^\perp\| \leq b_0$, $\|(\mathbf{x}^s, \mathbf{u}^s)\| \leq \gamma_0 \|\mathbf{b}^\perp\|$. There exists $\epsilon > 0$ such that for all $\|(\mathbf{x}(0), \mathbf{u}(0))\| < \epsilon$, $\|(\mathbf{x}(t) - \mathbf{x}^s, \mathbf{u}(t) - \mathbf{u}^s)\| \leq Ae^{-t/\tau}$ for some $A, \tau > 0$.*

C. *Suppose the region of influence of agent i_0 is the entire network and that $\bar{a}_{i_0 i_0} = 1$. For any $\nu > 0$, there exist thresholds $b_{th} > 0$ and $u_{th} > 0$ such that, if $|b_{i_0}^\perp| > b_{th}$*

and $u_f > u_{th}$, then there exists an equilibrium $(\mathbf{x}^s, \mathbf{u}^s)$ in the orthant $\{u_k \geq 0, \text{sign}(x_k) = \text{sign}(b_k^\perp), k = 1, \dots, N_a\}$, satisfying $|u_k^s - u_{sat}| < \nu$, for all $k \in \{1, \dots, N_a\}$, such that, $\lim_{t \rightarrow \infty} \|(\mathbf{x}(t) - \mathbf{x}^s, \mathbf{u}(t) - \mathbf{u}^s)\| = 0$.

Proof. The proofs of statements A, B are directly analogous to the proofs of Theorem VI.6-A, B and we leave out the details due to redundancy and space constraints.

C. Modulo a change of coordinate of the form $x_i \mapsto \pm x_i$, $i = 1, \dots, N_a$, we can assume $b_i^\perp > 0$ for all $i = 1, \dots, N_a$. It follows that there exists $\varepsilon > 0$ such that the set $\mathcal{U}^- = \{u_k \geq 0, x_k \geq -\varepsilon, k = 1, \dots, N_a\}$ is forward invariant for (27),(30). Moreover, any equilibrium in the forward invariant set \mathcal{U}^- is contained in the set $\mathcal{U}^+ = \{u_k \geq 0, x_k \geq +\varepsilon, k = 1, \dots, N_a\} \subset \mathcal{U}^-$. The flow generated by (27),(30) is monotone in \mathcal{U}^- and any trajectory converges to some equilibrium $(\mathbf{x}^s, \mathbf{u}^s)$. It remains to show that $|u_k^s - u_{sat}| < \nu$ for all $k = 1, \dots, N_a$, provided $b_{i_0}^\perp$ and u_f are large enough. Let i_2, \dots, i_{N_a} be the vertices along the influence path starting from i_0 , i.e., $\bar{a}_{i_2 i_0} = \bar{a}_{i_{k+1} i_k} = 1, k = 2, \dots, N_a - 1$.

The attention dynamics of the distinguished node i_0 reads $\dot{u}_{i_0} = -u_{i_0} + S_u(x_{i_0}^2 + \sum_{k \neq i_0} a_{i_0 k} x_k^2)$. If $a_{i_0 k} = 0$ for all $k \neq i_0$, then it follows by Theorem VI.6 that $x_{i_0}^s$ can be made arbitrary large and $u_{i_0}^s$ arbitrary close to u_{sat} by increasing the value of $b_{i_0}^\perp$. The same holds true in the general case, because $S_u(x_{i_0}^2 + \sum_{k \neq i_0} a_{i_0 k} x_k^2) \geq S_u(x_{i_0}^2)$.

The attention dynamics of the second node i_2 along the influence path originating from i_0 reads $\dot{u}_{i_2} = -u_{i_2} + S_u(x_{i_2}^2 + \sum_{k \neq i_2} a_{i_2 k} x_k^2)$. At steady-state, $u_{i_2}^s = S_u((x_{i_0}^s)^2 + \sum_{k \neq i_0} a_{i_0 k} (x_k^s)^2)$. Because $x_{i_0}^s$ is arbitrary large for sufficiently large $b_{i_0}^\perp$, it follows that $u_{i_2}^s$ is arbitrarily close to u_{sat} for sufficiently large $b_{i_0}^\perp$. Recall that $u_{sat} = u_f e^{g y_m} / (1 + e^{g y_m})$, thus u_{sat} can be made arbitrary large by increasing the value of u_f . In turns, by implicit differentiation of the steady-state equation $0 = -x_{i_2}^s + u_{i_2}^s \hat{S}(x_{i_2}^s) + b_{i_2}^\perp$, it follows that $\frac{\partial x_{i_2}^s}{\partial u_{i_2}^s} \geq c > 0$ for all $(x_{i_2}^s, u_{i_2}^s) \in \mathcal{U}^+$. Thus, by increasing u_f , $x_{i_2}^s$ can also be made arbitrarily large. The rest of the proof follows by repeating the same construction for the remaining nodes i_3, \dots, i_{N_a} along the path originating from i_0 . \square

Fig. 10 illustrates a cascade as predicted by Theorem VI.8 for 5 agents with only agent 1 getting a large input. Because agent 1 is at the root of the path graph that defines the attention dynamic coupling, its response to its input sets off a cascade such that all the agents form a strong opinion in the direction of their small inputs. Without the influence of agent 1, the other agents would not have formed strong opinions.

The coexistence and multistability of consensus and dissensus equilibria, as well as other interpretable equilibria, make the decoupled opinion dynamics maximally flexible and sensitive to input. Theorem VI.8 shows that, with the addition of attention feedback dynamics, agents reliably favor the option informed by their input, and agents without input reliably flip a coin and form an opinion in favor of one of the options. Due to the bistability of various branches of equilibria near the singularity, a change in input can easily facilitate




Fig. 10: Simulation of dynamics (27),(30) show a cascade for $N_a = 5$ agents from initial condition $(x_i(0), u_i(0)) = (0, 0)$, for all i and agent 1 the only agent getting a large input. The attention dynamics self weights are $\bar{a}_{ii} = 1$ and nonzero inter-agent weights are give by the directed path graph: $\bar{a}_{21} = \bar{a}_{32} = \bar{a}_{43} = \bar{a}_{54} = 1$. $\alpha = 1, \beta = -1, u_f = 2, g = 10, y_m = 1.5, \tau_u = 1, \mathbf{b} = (1.5, -0.015, -0.1524, 0.201, 0.0011)$.

a transition from one configuration of opinions to another, including transitions between consensus and dissensus. The design parameters u_f, y_m can be used to regulate the sensitivity threshold and robustness threshold of the group's opinion cascade in the same way outlined for uncoupled agents in Section VI-B.

D. Tunable Transitions between Consensus and Dissensus

In this section we illustrate how introducing feedback dynamics to social influence weights in the opinion dynamics can be used to facilitate transitions between group consensus and dissensus opinion configurations.

We focus on an all-to-all network with $N_o = 2$ options and N_a agents with interaction weights that satisfy the conditions in Theorem III.6 for reduction of opinion dynamics (20) to opinion dynamics of $N_c = 2$ clusters. We consider parameters as follows:

- 1) $u_i = u, d_i = d$ for all $i = 1, \dots, N_a$;
- 2) $b_i^\perp = b_p^\perp$ when $i \in \mathcal{I}_p, p \in \{1, 2\}$;
- 3) $A_{ik}^{11} = A_{ik}^{22} = \alpha/N_p, A_{ik}^{12} = A_{ik}^{21} = \beta/N_p$, when $i, k \in \mathcal{I}_p, p \in \{1, 2\}$;
- 4) $A_{ik}^{11} = A_{ik}^{22} = \gamma_{ik}/N_s, A_{ik}^{12} = A_{ik}^{21} = \delta_{ik}/N_s$ when $i \in \mathcal{I}_p, k \in \mathcal{I}_s, s, p \in \{1, 2\}$ and $s \neq p$.

The opinions of the group converge to a two-dimensional manifold \mathcal{E} on which the effective $(S_2 \times S_2)$ -equivariant opinion dynamics are

$$\dot{\hat{x}}_p = -d\hat{x}_p + u \left(\hat{S}_1(\alpha\hat{x}_p + \gamma\hat{x}_s) - \hat{S}_2(\beta\hat{x}_p + \delta\hat{x}_s) \right) + b_p^\perp \quad (33)$$

with $p, s \in \{1, 2\}, p \neq s$, where \hat{x}_p is the average opinion of cluster p :

$$\hat{x}_p = \frac{1}{N_p} \sum_{i \in \mathcal{I}_p} x_i, \quad p \in \{1, 2\}. \quad (34)$$

We apply the dynamic feedback for attention (30) and design distributed state feedback dynamics for the inter-agent inter-

action weights γ_i, δ_i , where each agent i in cluster p has an opinion that evolves according to

$$\dot{x}_i = -dx_i + u \left(\hat{S}_1(\alpha \hat{x}_p + \gamma_i \hat{x}_s) - \hat{S}_2(\beta \hat{x}_p + \delta_i \hat{x}_s) \right) + b_i^\perp \quad (35)$$

with $p, s \in \{1, 2\}$, $p \neq s$. These dynamics also take the form of a leaky nonlinear integrator:

$$\tau_\gamma \dot{\gamma}_i = -\gamma_i + \sigma S_\gamma(\hat{x}_1 \hat{x}_2) \quad (36a)$$

$$\tau_\delta \dot{\delta}_i = -\delta_i - \sigma S_\delta(\hat{x}_1 \hat{x}_2) \quad (36b)$$

where $\sigma \in \{1, -1\}$, $\tau_\gamma, \tau_\delta > 0$ are time scales, and the saturating function is

$$S_c(y) = c_f \tanh(g_c y), \quad c \in \{\gamma, \delta\} \quad (37)$$

where the parameters $c_f, g_c > 0$.

Remark VI.5. *When all agents have the same initial condition for the interaction weights $\gamma_i(0), \delta_i(0)$, then $\gamma_i(t) = \gamma_k(t)$ and $\delta_i(t) = \delta_k(t)$ for all time and all $i, k \in \{1, \dots, N_a\}$.*

Any configuration of opinions in which \hat{x}_1 and \hat{x}_2 are non-neutral will drive γ_i to approach $\sigma\gamma_f$ and δ_i to approach $-\sigma\delta_f$. When $\sigma = 1$ this final state corresponds to a clustered consensus state and when $\sigma = -1$ it corresponds to a clustered dissensus state. Thus, introducing feedback dynamics to the opinion network weights can lead to formation of consensus or dissensus. The sign of design parameter σ determines whether the system tends towards consensus or dissensus, and switching the sign can reliably trigger a transition from consensus to dissensus and vice versa. Fig. 11 illustrates the opinion formation of 7 agents that form two clusters, one with 3 agents and the other with 4 agents. Initially, $\gamma - \delta < 0$, and the clusters evolve to a dissensus state. However, because $\sigma = 1$, the two clusters evolve towards a consensus state once $\gamma - \delta > 0$. At time $t = 300$, design parameter $\sigma = -1$ and the two clusters evolve towards a dissensus state once $\gamma - \delta < 0$.

VII. FINAL REMARKS

We have proposed and analyzed a new general model of continuous-time opinion dynamics for an arbitrary number of agents that interact over a network as they form opinions about an arbitrary number of options. The new model generalizes a great many models of opinion dynamics in the literature. It also exhibits the wide range of opinion formation behaviors that are predicted by model-independent theory. This includes the rapid and reliable formation of both consensus and dissensus states, even for highly symmetric and homogeneous groups.

A special feature of the model is that the range of possible behaviors can be distinguished by a small number of parameters, and this lends analytical tractability as well as the opportunity for systematizing design. The opinion formation process is also robust to small heterogeneity in parameters, initial conditions, and inputs. So, for example, results on idealized, symmetric networks, hold up in the presence of small variations and uncertainties in parameters.

We have shown that multistability of opinion formation outcomes in the model is another special feature since it yields an opinion formation process that exhibits ultra-sensitivity




Fig. 11: Top) opinion trajectories; Bottom) parameter trajectories. Seven agents form two clusters of sizes $N_1 = 3$ (dashed-line opinion trajectories), $N_2 = 4$ (solid-line opinion trajectories). Parameters are $d = 1$, $\alpha = 1$, $\beta = -1$, $\hat{b}_1^\perp = 0.5$, $\hat{b}_2^\perp = -0.5$, $\tau_u = 10$, $\tau_\gamma = \tau_\delta = 100$, $\gamma_f = 2$, $\delta_f = 1$, $u_f = 2$, $g = g_c = 10$, $y_m = 1.5$. Initially, $\gamma - \delta < 0$ and the group converges to a clustered dissensus state. For $t < 300$ $\sigma = 1$ and the group switches to a consensus state after $\gamma - \delta$ becomes positive. For $t \geq 300$, $\sigma = -1$ and the group switches from consensus to dissensus after $\gamma - \delta$ changes sign back to negative.

to inputs, robustness to changes in inputs, opinion cascades, and flexible transitions between consensus and dissensus. We have proposed feedback dynamics for the parameter we refer to as the agent's attention to social influence and we show how design parameters can be used to tune the sensitivity, robustness, cascades, and transitions.

In ongoing work we are studying the explicit role of network structure in opinion formation and in the tuning of sensitivity, robustness, cascades, and transitions. We are exploring the opinion formation process as the number of options grows, and we are using the model to accommodate opinion formation in the case of multiple issues using multi-layer networks. We are applying the general model to design multi-robot coordination, task switching, and decision making for spatial navigation and to explore mechanisms that explain social behavior from animal group foraging to political polarization.

REFERENCES

- [1] M. H. DeGroot, "Reaching a consensus," *Journal of the American Statistical Association*, vol. 69, no. 345, pp. 121–132, 1974.
- [2] N. E. Friedkin and E. C. Johnsen, "Social influence networks and opinion change," in *Advances in Group Processes*, S. R. Thye, E. J. Lawler, M. W. Macy, and H. A. Walker, Eds. Emerald Group Publishing Limited, 1999, vol. 16, p. 129.
- [3] P. Cisneros-Velarde, K. S. Chan, and F. Bullo, "Polarization and fluctuations in signed social networks," *ArXiv*, vol. abs/1902.00658, 2019.
- [4] R. Olfati-Saber and R. M. Murray, "Consensus problems in networks of agents with switching topology and time-delays," *IEEE Transactions on Automatic Control*, vol. 49, no. 9, pp. 1520–1533, 2004.
- [5] C. Altafini, "Consensus problems on networks with antagonistic interactions," *IEEE Transactions on Automatic Control*, vol. 58, no. 4, pp. 935–946, 2013.
- [6] G. Deffuant, D. Neau, F. Amblard, and G. Weisbuch, "Mixing beliefs among interacting agents," *Advances in Complex Systems*, vol. 03, no. 01n04, pp. 87–98, 2000.

- [7] R. Hegselmann and U. Krause, "Opinion dynamics and bounded confidence models, analysis, and simulations," *Journal of Artificial Societies and Social Simulation*, vol. 5, no. 3, pp. 121–132, 2002.
- [8] —, "Opinion dynamics driven by various ways of averaging," *Computational Economics*, vol. 25, no. 4, pp. 381–405, 2005.
- [9] V. Blondel, J. M. Hendrickx, and J. N. Tsitsiklis, "On Krauses multi-agent consensus model with state-dependent connectivity," *IEEE Transactions on Automatic Control*, vol. 54, no. 11, pp. 2586–2597, 2009.
- [10] R. Sepulchre, "Consensus on nonlinear spaces," *IFAC Proceedings Volumes*, vol. 43, no. 14, pp. 1029 – 1039, 2010.
- [11] M. Caponigro, A. C. Lai, and B. Piccoli, "A nonlinear model of opinion formation on the sphere," *Discrete & Continuous Dynamical Systems - A*, vol. 35, no. 9, pp. 4241–4268, Sep. 2015.
- [12] L. Ding, W. X. Zheng, and G. Guo, "Network-based practical set consensus of multi-agent systems subject to input saturation," *Automatica*, vol. 89, pp. 316–324, 2018.
- [13] A. Hu, J. Cao, M. Hu, and L. Guo, "Event-triggered group consensus for multi-agent systems subject to input saturation," *Journal of the Franklin Institute*, vol. 355, no. 15, pp. 7384–7400, 2018.
- [14] P. Dandekar, A. Goel, and D. T. Lee, "Biased assimilation, homophily, and the dynamics of polarization," *Proceedings of the National Academy of Sciences*, vol. 110, no. 15, pp. 5791–5796, 2013.
- [15] A. Franci, V. Srivastava, and N. E. Leonard, "A realization theory for bio-inspired collective decision making," *arXiv:1503.08526 [math.OC]*, Mar. 2015.
- [16] R. Gray, A. Franci, V. Srivastava, and N. E. Leonard, "Multiagent decision-making dynamics inspired by honeybees," *IEEE Transactions on Control of Network Systems*, vol. 5, no. 2, pp. 793–806, 2018.
- [17] A. Fontan and C. Altafini, "Achieving a decision in antagonistic multi agent networks: frustration determines commitment strength*," in *IEEE Conference on Decision and Control*, 2018, pp. 109–114.
- [18] W. Xia, M. Ye, J. Liu, M. Cao, and X.-M. Sun, "Analysis of a nonlinear opinion dynamics model with biased assimilation," *arXiv:1912.01778 [math.OC]*, Dec. 2019.
- [19] B. Nabet, N. E. Leonard, I. D. Couzin, and S. A. Levin, "Dynamics of decision making in animal group motion," *Journal of Nonlinear Science*, vol. 19, pp. 399–435, 2009.
- [20] N. E. Leonard, T. Shen, B. Nabet, L. Scardovi, I. D. Couzin, and S. A. Levin, "Decision versus compromise for animal groups in motion," *Proceedings of the National Academy of Sciences*, vol. 109, no. 1, pp. 227–232, 2012.
- [21] D. Pais, P. M. Hogan, T. Schlegel, N. R. Franks, and N. E. Leonard, "A mechanism for value-sensitive decision-making," *PLoS One*, vol. 8, p. e73216, 2013.
- [22] I. Pinkoviezky, I. D. Couzin, and N. S. Gov, "Collective conflict resolution in groups on the move," *Phys. Rev. E*, vol. 97, p. 032304, 2018.
- [23] A. F. Siegenfeld and Y. Bar-Yam, "Negative representation and instability in democratic elections," *Nature Physics*, vol. 16, pp. 186–190, 2020.
- [24] S. E. Parsegov, A. V. Proskurnikov, R. Tempo, and N. E. Friedkin, "A new model of opinion dynamics for social actors with multiple interdependent attitudes and prejudices," in *IEEE Conference on Decision and Control*, 2015, pp. 3475–3480.
- [25] N. E. Friedkin, A. V. Proskurnikov, R. Tempo, and S. E. Parsegov, "Network science on belief system dynamics under logic constraints," *Science*, vol. 354, no. 6310, pp. 321–326, 2016.
- [26] S. E. Parsegov, A. V. Proskurnikov, R. Tempo, and N. E. Friedkin, "Novel multidimensional models of opinion dynamics in social networks," *IEEE Transactions on Automatic Control*, vol. 62, no. 5, pp. 2270–2285, 2017.
- [27] S. Fortunato, V. Latora, A. Pluchino, and A. Rapisarda, "Vector opinion dynamics in a bounded confidence consensus model," *International Journal of Modern Physics C*, vol. 16, p. 15351551, 2005.
- [28] A. Nedić and B. Touri, "Multi-dimensional Hegselmann-Krause dynamics," in *IEEE Conference on Decision and Control*, 2012, pp. 68–73.
- [29] L. Li, A. Scaglione, A. Swami, and Q. Zhao, "Consensus, polarization and clustering of opinions in social networks," *IEEE Journal on Selected Areas in Communications*, vol. 31, no. 5, pp. 1072–1083, 2013.
- [30] L. Pan, H. Shao, M. Mesbahi, Y. Xi, and D. Li, "Bipartite consensus on matrix-valued weighted networks," *IEEE Transactions on Circuits and Systems II: Express Briefs*, vol. 66, no. 8, pp. 1441–1445, 2019.
- [31] M. Ye, J. Liu, L. Wang, B. D. O. Anderson, and M. Cao, "Consensus and disagreement of heterogeneous belief systems in influence networks," *IEEE Transactions on Automatic Control*, pp. 1–1, 2019.
- [32] A. Sirbu, V. Loreto, V. D. P. Servedio, and F. Tria, "Opinion dynamics with disagreement and modulated information," *Journal of Statistical Physics*, vol. 151, no. 1, pp. 218–237, 2013.
- [33] A. Franci, M. Golubitsky, A. Bizyaeva, and N. E. Leonard, "A model-independent theory of consensus and dissensus decision making," *arXiv:1909.05765 [math.OC]*, Sep. 2020.
- [34] P. Reverdy, "Dynamical, value-based decision making among n options," *arXiv:2003.03874*, Mar. 2020.
- [35] A. Reina, J. A. R. Marshall, V. Trianni, and T. Bose, "Model of the best-of- n nest-site selection process in honeybees," *Phys. Rev. E*, vol. 95, p. 052411, May 2017. [Online]. Available: <https://link.aps.org/doi/10.1103/PhysRevE.95.052411>
- [36] M. Golubitsky and I. Stewart, *The Symmetry Perspective*, 1st ed., ser. Progress in Mathematics. New-York: Birkhäuser Basel, 2002, vol. 200.
- [37] M. Golubitsky and D. G. Schaeffer, *Singularities and Groups in Bifurcation Theory*, ser. Applied Mathematical Sciences. New York, NY: Springer-Verlag, 1985, vol. 51.
- [38] J. J. Hopfield, "Neural networks and physical systems with emergent collective computational abilities," *Proceedings of the National Academy of Sciences*, vol. 79, no. 8, pp. 2554–2558, 1982.
- [39] —, "Neurons with graded response have collective computational properties like those of two-state neurons," *Proceedings of the National Academy of Sciences*, vol. 81, no. 10, pp. 3088–3092, 1984.
- [40] Y. Nakamura, K. Torii, and T. Munakata, "Neural-network model composed of multidimensional spin neurons," *Physical Review E*, vol. 51, no. 2, pp. 1538–1546, 1995.
- [41] M. Usher and J. L. McClelland, "The time course of perceptual choice: The leaky, competing accumulator model," *Psychological Review*, vol. 108, no. 3, pp. 550–592, 2001.
- [42] R. Bogacz and K. Gurney, "The basal ganglia and cortex implement optimal decision making between alternative actions," *Neural Computation*, vol. 19, no. 2, pp. 442–477, 2007.
- [43] J. Ghaderi and R. Srikant, "Opinion dynamics in social networks with stubborn agents: Equilibrium and convergence rate," *Automatica*, vol. 50, pp. 3209–3215, 2014.
- [44] J. Lorenz, "Continuous opinion dynamics of multidimensional allocation problems under bounded confidence: More dimensions lead to better chances for consensus," in *Eur. J. Econ. Soc. Syst.*, vol. 19, 2006, pp. 213–227.
- [45] P. Jia, A. MirTabatabaei, N. E. Friedkin, and F. Bullo, "Opinion dynamics and the evolution of social power in influence networks," *SIAM Review*, vol. 57, no. 3, pp. 367–397, 2015.
- [46] H. Khalil, *Nonlinear systems*, 3rd ed. Pearson Education International Inc., 2000.
- [47] R. Sepulchre, D. A. Paley, and N. E. Leonard, "Stabilization of planar collective motion: All-to-all communication," *IEEE Transactions on Automatic Control*, vol. 52, no. 5, pp. 811–824, 2007.
- [48] —, "Stabilization of planar collective motion with limited communication," *IEEE Transactions on Automatic Control*, vol. 53, no. 3, pp. 706–719, 2008.
- [49] S. H. Strogatz, "From Kuramoto to Crawford: Exploring the onset of synchronization in populations of coupled oscillators," *Physica D: Nonlinear Phenomena*, vol. 143, no. 1–4, pp. 1–20, 2000.
- [50] M. Golubitsky, I. Stewart, and D. Schaeffer, *Singularities and Groups in Bifurcation Theory, Vol. II*, ser. Applied Mathematical Sciences. New York, NY: Springer-Verlag, 1988, vol. 69.
- [51] K. Wang and N. Michel, "Robustness and perturbation analysis of a class of nonlinear systems with applications to neural networks," *IEEE Transactions on Circuits and Systems I: Fundamental Theory and Applications*, vol. 41, no. 1, pp. 24–32, Jan 1994.
- [52] E. Salinas and P. Thier, "Gain modulation: A major computational principle of the central nervous system," *Neuron*, vol. 27, no. 1, pp. 15 – 21, 2000.
- [53] E. Brown, J. Gao, P. Holmes, R. Bogacz, M. Gilzenrat, and J. D. Cohen, "Simple neural networks that optimize decisions," *International Journal of Bifurcation and Chaos*, vol. 15, no. 03, pp. 803–826, 2005.
- [54] S. Tajima, J. Drugowitsch, N. Patel, and A. Pouget, "Optimal policy for multi-alternative decisions," *Nature Neuroscience*, vol. 22, no. 9, pp. 1503–1511, 2019.
- [55] Y. D. Zhong and N. E. Leonard, "A continuous threshold model of cascade dynamics," in *2019 IEEE 58th Conference on Decision and Control (CDC)*. IEEE, 2019, pp. 1704–1709.
- [56] H. L. Smith, *Monotone Dynamical Systems: An Introduction to the Theory of Competitive and Cooperative Systems*. American Mathematical Society, 2008, no. 41.