

Robust Stochastic Stability with Applications to Social Distancing in a Pandemic

Brandon C. Collins, Lisa Hines, Gia Barboza, and Philip N. Brown

Abstract—The theory of learning in games has extensively studied situations where agents respond dynamically to each other in light of a fixed utility function. However, in many settings of interest, agent utility functions themselves vary as a result of past agent choices. The ongoing COVID-19 pandemic has highlighted the need to formulate and analyze such models which feature game-environment feedback. For instance, a highly prevalent virus may incentivize individuals to wear masks, but extensive adoption of mask-wearing reduces virus prevalence which in turn reduces individual incentives for mask-wearing. What is the interplay between epidemic severity and the behaviors of a victim population? For initial answers, we develop a general framework using probabilistic coupling methods that can be used to derive the stochastically stable states of log-linear learning in certain games which feature such game-environment feedback. We then apply this framework to a simple dynamic game-theoretic model of social precautions in an epidemic and give conditions under which maximally-cautious social behavior in this model is stochastically stable.

I. INTRODUCTION

In social systems and distributed engineered systems, collective behavior is the result of many individuals making intertwined self-interested choices. In many cases, the value of a particular choice depends not only on the current choices being made by others, but also on the history of past choices.

The recent COVID-19 pandemic exemplifies this. In particular, many social conventions such as wearing masks and social distancing are known to be effective at reducing the contagiousness of the disease [1], [2]. To effectively deploy mitigation policies, it is critical to understand how a population’s willingness to adopt preventative conventions interacts dynamically with the severity of an epidemic. For example, in the absence of an epidemic a population may prefer not to practice social distancing, but rising case counts may incentivize individuals to change their behavior to avoid contracting the disease. However, an epidemic’s severity at any given time is not only a function of its own dynamics, but also the behavioral history of the victim population. The resulting feedback loop may lead to challenges in predicting the effectiveness of mitigation policies.

In principle, these socio-environmental feedback loops can be analyzed using techniques from *game theory*, which has a long history of analyzing the society-scale effects of self-interested behavior. For instance, game theory has long been used to study the spread of social conventions [3] using models such as the graphical coordination game [4]

with the stochastic learning algorithm log-linear learning [5]. However, traditional analysis techniques almost uniformly assume that the game’s utility functions are fixed for all time, so that the agents’ choices over time can be described by a stationary Markov process. However, such analysis fails or becomes unwieldy when utility functions themselves depend on the history of play.

Analysis techniques for history-dependent games have broad potential applications; for instance, game theoretic methods are frequently proposed in the area of distributed control of multiagent systems [6]–[9]. However, in a distributed control application, agents’ actions may directly modify the strategic environment; for instance if a search-and-rescue UAV identifies a disaster victim, that victim may be removed from the list of other UAVs’ objectives. Other applications that can be modeled by history-dependent games are in machine learning [10]–[12] and biology [13], [14].

Owing in part to the challenges of modeling the complex game-environment feedback inherent to history-dependent games, general results on these games are elusive. Recent work has focused on specific learning algorithms and strategic environments, such as zero-sum games under replicator dynamics [15]. In [16] the authors characterize an oscillating tragedy of the commons effect under certain environmental feedback scenarios.

In this paper, we develop a general framework for analyzing the long-run behavior of binary-action history-dependent games. In particular, we study the stochastically stable states of the popular log-linear learning algorithm in such settings. We show that if the utility functions of the history-dependent game can be appropriately referenced to the utility functions of a corresponding exact potential game, then the history-dependent game of interest inherits the stochastically stable states of the reference potential game. To accomplish this we apply techniques from the theory of probabilistic couplings, and derive a monotone coupling that relates play in the history-dependent game with that in the reference potential game. To showcase an application of the framework, we present an epidemic model that intertwines the compartmental SIS disease model with the graphical coordination game convention model. Using our analysis framework we provide conditions under which the stochastically stable states may be fully characterized, despite their history-dependence.

A. A Motivating Epidemic Example

The novel COVID-19 epidemic has led to a surge in interest in understanding social responses to epidemics [17]–[19]. To model such a scenario we adopt a game theoretic model of behavior intertwined with a compartmental disease

This work was supported by the National Science Foundation under Grants #DEB-2032465 and #ECCS-2013779.

The authors are with the University of Colorado at Colorado Springs, CO 80918, USA {bcollin3, lhines, gbarboza, philip.brown}@uccs.edu




Fig. 1. Two runs of the SISGCG model. The full description can be found in Section IV however a brief description of the parameters is as follows. The infectiousness of an agent using no precautions is β_0 and using precautions is β_1 . The curing parameter, determining how fast agents recover, is γ .

model. Specifically, we couple a graphical coordination game with a compartmental SIS epidemic model. The coordination game is a model for how social conventions spread in society [3], using the assumption that one gains intrinsic value to using the same conventions as others around them. In this model, which we call *SISGCG*, agents choose whether or not they take precautions by considering both a desire to coordinate on a set of social distancing conventions and a desire to practice safe conventions, influenced by the severity of the epidemic. The dynamics and thus severity of the epidemic are then in turn impacted by agent's decision to adopt precautions. We present the complete details of the SISGCG model in Section IV. The upper subplot of Figure 1 showcases how SISGCG captures the complex interplay between epidemic dynamics and social response. In particular, epidemic peaks in this model occur due to recent history of relatively many agents choosing not to take precautions. In addition, both social conventions and the epidemic take time to spread in SISGCG, so SISGCG models a complex relationship between delayed game theoretic social response, reactive epidemic dynamics, and the curing time of the disease.

The lower subplot of Figure 1 showcases a scenario in which our analysis framework can be used to characterize agent behaviour using stochastic stability. Note that the lower subplot depicts the case that social distancing is desirable even when the disease is at a lower-prevalence state and thus would also be desirable at any higher disease state.

II. MODEL

A. Game Formulation

In this work we consider binary action games. Letting $N = \{1, 2, 3, \dots, |N|\}$ denote the player set, in a binary action game player $i \in N$ has action set $A_i = \{0, 1\}$. The joint action space is then given by $A = \{0, 1\}^{|N|}$. We denote an action profile as $a \in A$ and use a_i to denote player i 's action and the actions of all other players by $a_{-i} = (a_1, a_2, \dots, a_{i-1}, a_{i+1}, \dots, a_{|N|})$. We refer to the all

1 action profile as $\vec{1} = (1)_{i=1}^{|N|}$ and similarly for the all zero profile, $\vec{0}$. Further, let $\Delta(A)$ denote the standard probability simplex over A .

Let $U_i : A \rightarrow \mathbb{R}$ be player i 's utility function. We denote $U = \{U_i\}_{i \in N}$ as the collection of all players utility function. Thus we may denote a game using tuple (N, A, U) , and let \mathcal{G} be the set of all such tuples.

A game $g \in \mathcal{G}$ is an *exact potential game* if there exists a potential function ϕ such that

$$U_i(a'_i, a_{-i}) - U_i(a_i, a_{-i}) = \phi(a'_i, a_{-i}) - \phi(a_i, a_{-i}) \quad (1)$$

for any $a \in A$, and $a_i, a'_i \in A_i$.

However, games $g \in \mathcal{G}$ cannot be used to model scenarios where play interacts with a dynamic environment. That is, the payoffs of such games cannot depend on the history of play that may impact the players' strategic environment. In this work we generalize \mathcal{G} by allowing each history of play to have a unique utility function. We write \mathcal{A}_T to denote the set of joint action histories of length $T \in \mathbb{N}$, and the set of all histories as $\mathcal{A} = \cup_{T \in \mathbb{N}} \mathcal{A}_T$. We typically use $\alpha \in \mathcal{A}_T$ to refer to a path and use superscripts to refer to time indices. For example, $\alpha^1 \in A$ is the first action profile in the history and α^T is the last, noting we use the convention that T is the last time index even when $\alpha \in \mathcal{A}$. We also define A, \mathcal{A}_T as partially ordered sets by first defining partial order \geq_A , where $a' \geq_A a$ whenever $a, a' \in A$ and $a'_i \geq a_i$ for all $i \in N$, recalling that $a'_i, a_i \in \{0, 1\}$. Using this we define partial order $\geq_{\mathcal{A}_T}$ as $\bar{\alpha} \geq \alpha$ whenever $\alpha, \bar{\alpha} \in \mathcal{A}_T$ and $\bar{\alpha}^t \geq_A \alpha^t$ for all $t \in \{1, 2, \dots, T\}$.

Using these notions we generalize the previously given game formulation \mathcal{G} to allow for the utility functions to depend on the history of play. Similar to before, let $U_i^\alpha : A \rightarrow \mathbb{R}$, where this utility function is not only specific player i but also to the history α . As before, let $U^\alpha = (U_1^\alpha, U_2^\alpha, \dots, U_{|N|}^\alpha)$ denote each player's utility function given history α and let $U^A = \{U^\alpha \mid \alpha \in \mathcal{A}\}$ be the set of utility functions across all paths. We denote a *history-dependent game* as tuple (N, A, U^A) and let \mathcal{G}^A be the set of all such tuples. We now present a class of games that combines potential games and history dependence.

Definition 1: We call a tuple $g = (N, A, U^A) \in \mathcal{G}^A$ an *aligned history-dependent game* if there exists an exact potential game $\hat{g} = (N, A, \hat{U}) \in \mathcal{G}$ with potential function $\hat{\phi}$ such that:

- 1) $\{\vec{1}\} = \arg \max_{z \in A} \hat{\phi}(z)$
- 2) $U_i^\alpha(1, \alpha_{-i}^T) \geq \hat{U}_i(1, a_{-i})$
- 3) $\hat{U}_i(0, a_{-i}) \geq U_i^\alpha(0, \alpha_{-i}^T)$

for any $\alpha, \alpha' \in \mathcal{A}$, $a, a' \in A$, $T \in \mathbb{N}$ such that $\alpha_{-i}^T \geq_{A_{-i}} a_{-i}$ and a, a' vary by only a unilateral deviation. For convenience we denote ordering $\geq_{A_{-i}}$ over $A_{-i} = \{0, 1\}^{|N|-1}$ equivalently to \geq_A .

B. Learning in Games

In this work we focus on the learning algorithm log-linear learning, which is a discrete time asynchronous learning algorithm [5], [20]. That is, for game $g \in \mathcal{G}^A$ at each time

step log-linear learning will select a single agent to update their action to a_i with probability

We define the probability that agent i selects action a_i given history $\alpha \in \mathcal{A}_T$ as

$$\mathbb{P}_i^\alpha(a_i) = \frac{e^{\frac{1}{\tau}U_i^\alpha(a_i, \alpha_{-i}^T)}}{\sum_{a'_i \in A_i} e^{\frac{1}{\tau}U_i^\alpha(a'_i, \alpha_{-i}^T)}}. \quad (2)$$

where τ , called the *temperature* is a parameter which governs the rationality of agents. As $\tau \rightarrow 0$ agents will best respond with high probability, and as $\tau \rightarrow \infty$ agents will choose actions uniformly at random. Note that we take the last action profile in the history α^T as the behavior of the other agents. We define the probability $\alpha \in \mathcal{A}_T$ transitions to $a' \in A$ under log-linear learning in a single transition as

$$P^\alpha(a') = \begin{cases} \frac{1}{|N|} \sum_{j \in N} \mathbb{P}_j^\alpha(a'_j) & \alpha^T = a' \\ \frac{1}{|N|} \mathbb{P}_i^\alpha(a'_i) & \alpha_i^T \neq a'_i, \alpha_{-i}^T = a'_{-i} \\ 0 & \text{else.} \end{cases} \quad (3)$$

This can be interpreted as the probability that given history α the next action profile $\alpha^{T+1} = a'$.

We say $a \in A$ is *strictly stochastically stable* if the following ϵ definition [21] holds. For any $\epsilon > 0$ there exists $T > 0, T < \infty$ such that such that

$\Pr(s(t; \tau, \pi, g) = \vec{1}) > 1 - \epsilon$ whenever $t > T, \tau < T$. (4) where $s(\cdot)$ is a random variable representing the action profile at time t under log-linear learning, given temperature τ , initial distribution $\pi \in \Delta(A)$ and game g .

Traditionally, exact potential games under log-linear learning may be analyzed using a theory of *resistance trees* [3], [5], [20], [22] to relate potential function maximizers to stochastic stability. However, this analysis depends on the fact that log-linear learning induces an ergodic markov chain from any exact potential game. It is easy to show that when log-linear learning is applied to game $g \in \mathcal{G}^A$ the induced random process is not Markovian in general. In particular, the Markov property requires that at any time determining the next state does not depend on the past, however the underlying utility functions of g explicitly depend on history α , and therefore the associated transition given in (2) violates the Markov property. Thus the traditional technique for showing stochastic stability in this context fails.

III. MAIN CONTRIBUTION

We now present our main result, extending stochastic stability to the class of games we term aligned history-dependent games. In particular, we show that for such games the $\vec{1}$ state is strictly stochastically stable.

Theorem 1: If $g \in \mathcal{G}^A$ is an aligned history-dependent game then $\vec{1}$ is strictly stochastically stable in g under log-linear learning.

The proof of Theorem 1 proceeds using Lemma 1, which we present here and prove in Section V. The interpretation of this lemma is that for an aligned history dependent game g , the probability at any time step that g is in the $\vec{1}$ action profile is lower bounded by the probability its associated exact potential game \hat{g} is in the $\vec{1}$ profile.

Lemma 1: If $g \in \mathcal{G}^A$ is an aligned history-dependent game with associated exact potential game $\hat{g} \in \mathcal{G}$ then

$\Pr(s(T; \tau, \pi, g) = \vec{1}) \geq \Pr(s(T; \tau, \pi, \hat{g}) = \vec{1})$ for any temperature $\tau > 0, \pi \in \Delta(A)$.

The proof of Lemma 1 is technically involved and depends on our novel monotone coupling framework which we present in Section V. Using this result we now present a straightforward proof of Theorem 1.

Proof of Theorem 1: Let $g \in \mathcal{G}^A$ be an aligned history-dependent game and \hat{g} be its associated exact potential game.

It is well-known [20] that in an exact potential game $a \in A$ is a stochastically stable state under log-linear learning if

$$a \in \arg \max_{a' \in A} \phi(a'). \quad (5)$$

Therefore, because $\vec{1}$ is the lone maximizer of $\hat{\phi}$, it is strictly stochastically stable. We apply Lemma 1 directly to the definition of strict stochastic stability in (4). For any $\epsilon > 0$ there exists $T > 0, T < \infty$ such that such that

$$\Pr(s(t; \tau, \pi, g)) \geq \Pr(s(t; \tau, \pi, \hat{g})) > 1 - \epsilon \quad \forall t > T, \tau < T \quad (6)$$

yielding stochastic stability of $\vec{1}$ in game g . ■

IV. APPLICATION TO SOCIAL DISTANCING IN A PANDEMIC

A. SIS Preliminaries

It is appropriate to model epidemics that do not grant long term immunity with the SIS compartmentalized model. Here, we assume that every member of the population is in one of two states, susceptible (S) or infected (I). Individuals transition from susceptible to infected with rate $\beta > 0$, and infected to susceptible with rate $\gamma > 0$. The fraction $s(t)$ of the population that is susceptible at time t evolves as a function of time according to the one-dimensional nonlinear ODE

$$\dot{s} = -\beta s(1-s) + \gamma(1-s). \quad (7)$$

The solutions to (7) when $s(0) \in [0, 1]$ are fully characterized; in particular, it is known that when $\beta/\gamma > 1$, there exists an asymptotically stable equilibrium (called the *endemic equilibrium*) $s^* = \gamma/\beta$, and that every solution to (7) with $s(0) < 1$ converges to s^* . For more information, the reader may reference the survey [23].

B. The Graphical Coordination Game

One model for the spread of conventions in society is the Graphical Coordination Game (GCG) [4]. In this networked game, each agent plays the following two-player coordination game with each of her neighbors:

$$u(a_i, a_j) = 1 \begin{array}{|c|c|} \hline & \begin{array}{c} 1 \\ 0 \end{array} \\ \hline \begin{array}{c} q+c, q+c \\ 0, 0 \end{array} & \begin{array}{c} 0, 0 \\ 1, 1 \end{array} \\ \hline \end{array} \quad (8)$$

where $a_i \in \{0, 1\}$ is taken to be the row player's action and a_j is the column player's action, constant $q \in (0, 1]$ is a base coordination incentive and $c \geq 0$ is the *payoff gain*, indicating the benefit of coordinating on 1 over 0. The graphical coordination game is played between $|N|$ players on an undirected graph $G = (N, E)$ where E is the edge set. We define the neighbor set of any agent i as $\mathcal{N}_i = \{j \in N \mid (i, j) \in E, i \neq j\}$. In the GCG an agent's utility function for action profile $a \in A$ is given by

$$u_i(a) = \sum_{j \in \mathcal{N}_i} u(a_i, a_j). \quad (9)$$

When c is a constant it is well understood that the GCG in this formulation denotes an exact potential game. For convenience let $E_1(a) \subseteq E$ denote the set of edges whose agents are coordinated on action 1, and likewise for $E_0(a)$. The potential function is then given by

$$\phi(a) = (1+c)|E_1(a)| + |E_0(a)|. \quad (10)$$

C. A Socially Aware Epidemic Model

This model which we call SISGCG is defined by the following nonlinear hybrid dynamical system:

$$\dot{s} = (1-s)(\gamma - \beta(t)s). \quad (11)$$

Here, $\beta(t)$ is piecewise-constant and defined as follows: Let $\{t_k\}_{k \in \mathbb{N}} \in [0, \infty)$ be an unbounded set of agent revision opportunities. At each time t_k , an agent is selected uniformly at random to update her action, which she does according to log-linear learning with probability (2) with utilities associated with the following graphical coordination game, where we write $I(t) := 1 - s(t)$, and write player i 's action at time t as a_i^t :

$$u^{\text{SISGCG}}(a_i^t, a_j^t) = u(a_i^t, a_j^t) \text{ with } c = I(t) \quad (12)$$

where $q \in (0, 1]$ and represents how willing a population is to practice safe conventions in the absence of an epidemic. The action 1 represents a ‘‘safe convention’’ action in which a player is acting to reduce contagion; the action 0 represents conventions ignoring the pandemic. These actions are associated with infection coefficients $0 < \beta_1 < \beta_0$, respectively. Note in the SISGCG model we define the coordination game with log-linear learning in the same way as previously stated, but the 2-player coordination game is played with u^{SISGCG} in place of u . Finally, $\beta(t)$ is simply the average infection rate of all individuals, given their choices:

$$\beta(t) = \frac{1}{N} \sum_{i \in N} a_i^t \beta_1 + (1 - a_i^t) \beta_0. \quad (13)$$

Proposition 1: If $s(0) \in [0, 1)$, then if $s(t)$ is a solution of (11) with $\beta(t)$ given by (13), there exists a \bar{t} such that $s(t) \leq \gamma/\beta_1$ for all $t \geq \bar{t}$ almost surely.

Proof: We write $s_1^* := \gamma/\beta_1$. Note that if $s(t) \geq s_1^*$, then because $\beta(t) \geq \beta_1$, we have that $\dot{s} \leq 0$ by (11), and that this inequality is strict whenever $s(t) > s_1^*$. Thus, the set $[0, s_1^*]$ is positively invariant for the hybrid nonlinear dynamics given in (11).

To see that $s(t)$ eventually enters $[0, s_1^*]$ almost surely, consider the event that $s(t) > s_1^*$ for all t . Since s_1^* is asymptotically stable when $\beta(t) \equiv \beta_1$ and for any action profile $a \neq \vec{1}$ that its associated $\beta(t) > \beta_1$, it follows that the event that $\beta(t) \equiv \beta_1$ for all t is the same event as $s(0) > s_1^*$ and $s(t) > s_1^*$ for all t . However, it can be seen that the log-linear learning (3) action update probabilities define a stochastic process which visits every action profile in A infinitely often. That is, the probability that $\beta(t) \equiv \beta_1$ is 0, and thus there must exist a \bar{t} such that $s(t) \leq s_1^*$ for all $t \geq \bar{t}$ almost surely. ■

Immediately, it can be seen that this model exhibits some desirable intuitions about how populations behave both

during and in the absence of a pandemic. Consider the case with no pandemic present, that is $I(0) = 0$ which is a fixed point of the SIS dynamical system. In this scenario it can be seen that (12) is a constant payoff matrix, therefore this instance of SISGCG denotes an exact potential game. It is easy to see that $\vec{0}$ uniquely maximizes the potential function and therefore is strictly stochastically stable.

However, as a consequence of Proposition 1, if $I(0) > 0$ then $I(t) > 0$ for all t . It can be seen from (12) that SISGCG can be represented by a history-dependent game, as the utility function depends on the history of play. Therefore, agent decision-making inherits this dependence on the history of play, leading to non-Markovian behavior which complicates traditional analysis techniques. However, our Theorem 1 allows us to reference SISGCG to a related exact potential game and deduce conditions guaranteeing that $\vec{1}$ is strictly stochastically stable.

Proposition 2: Let g^S be an instance of SISGCG. If $\beta_1/\gamma > 1$, $q + \gamma/\beta_1 > 1$ and $I(0) > 0$ then we have $\vec{1}$ is stochastically stable in g .

Proof: To show stochastic stability of $\vec{1}$ in the SISGCG model we must:

- 1) show SISGCG is a history-dependent game,
- 2) show a corresponding fixed game \hat{g} exists, and
- 3) apply Theorem 1.

Let a SISGCG g^S be played on graph $G = (V, E)$ with $q + \gamma/\beta_1 > 1$ and $I(0) > 0$, and we consider g^S as played after time \bar{t} that exists almost surely as shown in Proposition 1. It is easy to show SISGCG is a history-dependent game, $g^S = (N, A, \bar{U})$. Note that trivially both the player set N and action set A match the history-dependent framework. To see that the utility functions fit into the framework note that the utility function, (12), depends on $I(t)$ which represents the proportion of agents who are infected. To find this constant for some $t \in \mathbb{N}$, it can be seen from (11) we require the history of agent up until that point. Thus we have at time t the utility functions \bar{U}^α depend on $\alpha \in \mathcal{A}_t$, which admits readily into the history-depend framework. Thus we have $g^S = (N, A, \bar{U}) \in \mathcal{G}^A$

Now we let $\hat{g}^S = (N, A, \hat{U}^S) \in \mathcal{G}$ be a GCG played on graph G . We define the pairwise coordination game payoff matrix by (8) with $c = \gamma/\beta_1$ inducing that \hat{g}^S denotes an exact potential game.

We now use \hat{g}^S to show g^S is an aligned history-dependent game. Because $q + \gamma/\beta_1 > 1$ its easy to see that $\vec{1}$ is the lone maximizer of the the potential function. Now we verify $U_i^\alpha(1, \alpha_{-i}^T) \geq \hat{U}_i^S(1, a_{-i})$ anytime $\alpha_{-i}^T \geq_{A_{-i}} a_{-i}$, $t > \bar{t}$. This can be rewritten for $t > \bar{t}$ as

$$\sum_{j \in \mathcal{N}_i^1(\alpha_{-i}^T)} q + I(t) \geq \sum_{j \in \mathcal{N}_i^1(a_{-i})} q + \gamma/\beta_1 \quad (14)$$

where $\mathcal{N}_i^1(a_{-i})$ denotes the neighbors of i who are playing 1 given profile a . This expressions holds because $\alpha_{-i}^T \geq_{A_{-i}} a_{-i} \Rightarrow |\mathcal{N}_i^1(\alpha_{-i}^T)| \geq |\mathcal{N}_i^1(a_{-i})|$ and by Proposition 1 $I(t) \geq \gamma/\beta_1$ for all $t > \bar{t}$. An argument with the same structure holds for $U_i^\alpha(0, \alpha_{-i}^T) \leq \hat{U}_i^S(0, a_{-i})$. Thus g^S is an aligned history-dependent game and we apply Theorem 1 to discover stochastic stability of $\vec{1}$ as desired. ■

V. PROOF VIA MONOTONE COUPLINGS

A. A Primer on Monotone Couplings

We begin with the definition of a monotone coupling, the core analytical device for our paper. Using this we present the definition of monotone coupling:

Definition 2: Let X be a countable set with partial ordering \leq_X and p_1, p_2 be probability measures on measure space (X, \mathcal{F}) . Then a *monotone coupling* of p_1, p_2 is a probability measure p on (X^2, \mathcal{F}^2) satisfying the following for all $x, y \in X$

$$\sum_{x \leq_X y'} p(x, y') = p_2(y') \text{ and } \sum_{y \geq_X x'} p(x', y) = p_1(x'). \quad (15)$$

A monotone coupling is a useful tool for analysis of the component probability measures p_1 and p_2 . In particular the following property holds in general for monotone couplings.

Proposition 3 (Paarporn et al., [24]): Let p_1, p_2 be probability measures on (X, \mathcal{F}) . If p is a monotone coupling of p_1, p_2 then for any increasing random variable $Z : X \rightarrow \mathbb{Z}_+$ we have

$$\mathbb{E}_{p_1}(Z) - \mathbb{E}_{p_2}(Z) = \sum_{\eta=0}^{\infty} p(Z_{\eta}^c, Z_{\eta}) \quad (16)$$

where $Z_{\eta} = \{a \mid Z(a) > \eta\}$.

Where we denote a complement set of $Z \subset X$ as Z^c . The proof is given in [24, Proposition 1].

B. Notation Required for Proofs

Taking $\hat{g} \in \mathcal{G}$, we give equations analogous to (2), (3) that give the transition probabilities for \hat{g} under log-linear learning. In particular, if agent i is selected to update her action then she will do so according to:

$$\hat{\mathbb{P}}_i^a(a_i) = \frac{e^{\frac{1}{\tau} U_i(a_i, a_{-i})}}{\sum_{a'_i \in A_i} e^{\frac{1}{\tau} U_i(a'_i, a_{-i})}} \quad (17)$$

Building on (17), we define the probability that action profile a transitions to a' under log-linear learning in a single transition as

$$\hat{P}^a(a') = \begin{cases} \frac{1}{|N|} \sum_{j \in N} \hat{\mathbb{P}}_j^a(a_j) & a = a' \\ \frac{1}{|N|} \hat{\mathbb{P}}_i^a(a'_i) & a_i \neq a'_i, a_{-i} = a'_{-i} \\ 0 & \text{else} \end{cases} \quad (18)$$

for some $i \in N$ and $a, a' \in A$. Additionally, we define the probability that path $\alpha \in \mathcal{A}_T$ occurs with initial distribution $\pi \in \Delta(A)$ as

$$\hat{P}_{\pi}(\alpha) = \pi(\alpha^1) \prod_{t=1}^{T-1} \hat{P}^{\alpha^t}(\alpha^{t+1}) \quad (19)$$

noting that $\pi(\alpha^1)$ denotes the probability of α^1 in initial distribution π .

Correspondingly, the probability that path $\alpha \in \mathcal{A}_T$ occurs with initial distribution $\pi \in \Delta(A)$ on $g \in \mathcal{G}^A$ is

$$P_{\pi}(\alpha) = \pi(\alpha^1) \prod_{t=1}^{T-1} P^{\alpha^{\leq t}}(\alpha^{t+1}) \quad (20)$$

where we use $\alpha^{\leq t} \in \mathcal{A}_t$ to mean history α until time $t \in \{1, 2, 3, \dots, T\}$.

We now present a result connecting the utility conditions of aligned history-varying potential games to (17) and (2).

Lemma 2: Let $g = (N, A, U^A) \in \mathcal{G}$, $\hat{g} = (N, A, \hat{U}) \in \mathcal{G}$ and let $i \in N, a \in A, \alpha \in \mathcal{A}$ such that $\alpha_{-i}^T \geq_{A_{-i}} a_{-i}$. If $U_i^{\alpha}(1, \alpha_{-i}^T) \geq \hat{U}_i(1, a_{-i})$ and $\hat{U}_i(0, a_{-i}) \geq U_i^{\alpha}(0, \alpha_{-i}^T)$ then $\mathbb{P}_i^{\alpha}(1) \geq \hat{\mathbb{P}}_i^a(1)$.

Proof: Let $g = (N, A, U^A) \in \mathcal{G}$, $\hat{g} = (N, A, \hat{U}) \in \mathcal{G}$ and let $i \in N, a \in A, \alpha \in \mathcal{A}$ be such that $\alpha_{-i}^T \geq_{A_{-i}} a_{-i}$. Further let $U_i^{\alpha}(1, \alpha_{-i}^T) \geq \hat{U}_i(1, a_{-i})$ and $\hat{U}_i(0, a_{-i}) \geq U_i^{\alpha}(0, \alpha_{-i}^T)$. Recalling $\tau > 0$, we begin by considering \mathbb{P}_i^{α}

$$\begin{aligned} \mathbb{P}_i^{\alpha}(1) &= \frac{e^{\frac{1}{\tau} U_i^{\alpha}(1, \alpha_{-i}^T)}}{e^{\frac{1}{\tau} U_i^{\alpha}(1, \alpha_{-i}^T)} + e^{\frac{1}{\tau} U_i^{\alpha}(0, \alpha_{-i}^T)}} \\ &\geq \frac{e^{\frac{1}{\tau} \hat{U}_i(1, a_{-i})}}{e^{\frac{1}{\tau} \hat{U}_i(1, a_{-i})} + e^{\frac{1}{\tau} \hat{U}_i(0, a_{-i})}} = \hat{\mathbb{P}}_i^a(1). \end{aligned} \quad (21)$$

To see the inequality, it suffices to apply the hypothesis to the fact that e^x and $l(x) = \frac{e^x}{e^x + c}$ are both increasing in x for $c > 0$. Thus $\mathbb{P}_i^{\alpha}(1) \geq \hat{\mathbb{P}}_i^a(1)$ holds as desired. ■

Our framework requires a careful partitioning of the action space into several sets corresponding to different types of agent action deviations. We use these formalizations for the various cases of the monotone coupling given in Theorem 2 as seen in (23).

Let $f : A \rightarrow 2^A$ be defined as $f(a) = \{a' \in A \mid a_i \neq a'_i, a_{-i} = a'_{-i} \text{ for } i \in N\}$ be the set of action profiles reachable from a via exactly one unilateral deviation. For $a, a' \in A$ let

$$g(a, a') = \begin{cases} i & a_i \neq a'_i \\ 0 & a = a' \\ -1 & a_i \neq a'_i, a_j \neq a'_j, i \neq j \end{cases} \quad (22)$$

indicate which agent unilaterally deviated their action between action profiles a, a' , and be -1 if multiple agents have deviated.

Now, let $a, a' \in A$ where $a' \geq_A a$. We denote several disjoint subsets of $f(a)$:

- 1) $r(a) = \{z \in f(a) \mid a_{g(a,z)} = 1\}$,
- 2) $q(a, a') = \{z \in f(a) \mid z \leq_A a'\} \setminus r(a)$, and
- 3) $s(a, a') = f(a) \setminus (q(a, a') \cup r(a))$.

These sets can be interpreted in the following way. The set $r(a)$ is the set of action profiles which decreased with respect to \geq_A and $q(\cdot)$, $s(\cdot)$ both increased. Between $q(\cdot)$ and $s(\cdot)$, $q(\cdot)$'s action profiles remain less than a' and $s(\cdot)$'s profiles are greater then or incomparable to a' . We now present three more analogous sets that are disjoint subsets of $f(a')$:

- 1) $R(a') = \{z \in f(a') \mid a'_{g(a',z)} = 0\}$,
- 2) $Q(a, a') = \{z \in f(a') \mid z \geq_A a\} \setminus R(a')$, and
- 3) $S(a, a') = f(a') \setminus (Q(a, a') \cup R(a'))$.

The interpretation of these sets are flipped relative to $r(\cdot)$, $q(\cdot)$ and $s(\cdot)$.

We now highlight some useful features of these sets. By their definitions it is evident that $q(\cdot), r(\cdot), s(\cdot)$ are a disjoint partition of $f(a)$, and that $Q(\cdot), R(\cdot), S(\cdot)$ are a disjoint partition of $f(a')$. For any $a, a', a' \geq_A a$ we relate these sets by a function $b^{a, a'} : f(a) \rightarrow f(a')$. To evaluate $b^{a, a'}(\bar{a})$, identify the agent who deviated their action between a, \bar{a} and then deviate that agent's action in a' . Formally, $b^{a, a'}(\bar{a}) =$

$(\neg a'_{g(a,\bar{a})}, a'_{-g(a,\bar{a})})$ where for convenience we define $\neg a_i \in \{0, 1\} \setminus \{a_i\}$ for $a_i \in A_i = \{0, 1\}$. In particular, this function relates the disjoint subsets of $f(a), f(a')$ according to the following lemma.

Lemma 3: If $a, a' \in A$ and $a \leq_A a'$, then the following statements hold:

- 1) $b^{a,a'} : r(a) \rightarrow S(a, a')$ is a bijection,
- 2) $b^{a,a'} : s(a, a') \rightarrow R(a')$ is a bijection, and
- 3) $b^{a,a'} : q(a, a') \rightarrow Q(a, a')$ is a bijection.

Lemma 3 is proved in the Appendix.

C. The One-Step Couplings

To prove Lemma 1 and obtain Theorem 1, we construct a monotone coupling $\nu_{\hat{g}}^a$ between measures P_{π}, \hat{P}_{π} . Noting that $\nu_{\hat{g}}^a$ is a coupling of measures over histories, we first construct a family of monotone couplings for each one-step transition (Theorem 2), and subsequently use these couplings to build the desired coupling over histories (Theorem 3).

Theorem 2: Let $g \in \mathcal{G}^A$ denote an aligned history-dependent game and $\hat{g} \in \mathcal{G}$ be its associated exact potential game. Then a monotone coupling exists between \hat{P}^a and P^{α} for any $\alpha \in \mathcal{A}, a \in A$ whenever $a \leq_A \alpha^T$. This monotone coupling $\nu^{a,\alpha} : A^2 \rightarrow [0, 1]$ is given in (23) in Figure 2.

Proof: Let $a \in A, \alpha \in \mathcal{A}$ such that $a \leq_A \alpha^T$ and let $g \in \mathcal{G}^A$ be an aligned history-dependent game where $\hat{g} \in \mathcal{G}$ is its associated exact potential game. To verify $\nu^{a,\alpha}$ is a monotone coupling we must show the following conditions from Definition 2 for any $\bar{a}, \bar{a}' \in A$:

- 1) $\nu^{a,\alpha}$ is a well-defined probability measure,
- 2) $\sum_{z' \geq_A \bar{a}} \nu^{a,\alpha}(\bar{a}, z') = \hat{P}^a(\bar{a})$, and
- 3) $\sum_{\substack{z' \geq_A \bar{a} \\ z \leq_A \bar{a}'}} \nu^{a,\alpha}(z, \bar{a}') = P^{\alpha}(\bar{a}')$.

We begin by verifying Condition 2. We consider cases $\bar{a} \notin (f(a) \cup \{a\}), \bar{a} \in q, \bar{a} \in r, \bar{a} \in s$ and $\bar{a} = a$ separately. We use the notational convention that q, s, Q, S are assumed to take arguments (a, α^T) and r, R take the argument a, α^T respectively. The first case represents any \bar{a} that cannot be achieved in a single unilateral deviation from a . Trivially, this gives that $\hat{P}^a(\bar{a}) = 0$, and thus all pairs of \bar{a}, z' must satisfy $\nu^{a,\alpha}(\bar{a}, z') = 0$. This holds as all parts of (23) require $\bar{a} \in (f(a) \cup \{a\})$ except (23h), which has the desired property.

We now consider the second case that $\bar{a} \in q$. Note that only (23d) satisfies this condition, so

$$\begin{aligned} \sum_{z' \geq_A \bar{a}'} \nu^{a,\alpha}(\bar{a}, z') &= \nu^{a,\alpha}(\bar{a}, \alpha^T) \\ &= \frac{\hat{P}_{g(a,\bar{a})}^a(\bar{a}_{g(a,\bar{a})})}{|N|} = \hat{P}^a(\bar{a}) \end{aligned} \quad (24)$$

as desired.

Next we consider $\bar{a} \in r$ which satisfies (23c), (23f) uniquely since b^{a,α^T} is a bijection by Lemma 3. Thus

$$\begin{aligned} \sum_{z' \geq_A \bar{a}'} \nu^{a,\alpha}(\bar{a}, z') &= \frac{1}{|N|} (\hat{P}_{g(a,\bar{a})}^a(0) \\ &\quad - \mathbb{P}_{g(a,\bar{a})}^{\alpha}(0) + \mathbb{P}_{g(\alpha^T, \bar{a}')}^{\alpha}(0)) \\ &= \frac{1}{|N|} \hat{P}_{g(a,\bar{a})}^a(0) = \hat{P}^a(\bar{a}) \end{aligned} \quad (25)$$

where the second equality follows as $g(a, \bar{a}) = g(\alpha^T, \bar{a}')$ by definition of b^{a,α^T} . The third equality follows as $\bar{a} \in r \implies \bar{a}_{g(a,\bar{a})} = 0$.

Considering $\bar{a} \in s$, we find only (23e) applies, thus for

$$\begin{aligned} \sum_{z' \geq_A \bar{a}'} \nu^{a,\alpha}(\bar{a}, z') &= \nu^{a,\alpha}(\bar{a}, b^{a,\alpha^T}(\bar{a})) \\ &= \frac{1}{|N|} \hat{P}_{g(a,\bar{a})}^a(1) = \hat{P}^a(\bar{a}) \end{aligned} \quad (26)$$

where $\bar{a} \in s \implies \bar{a}_{g(a,\bar{a})} = 1$ or else \bar{a} would be in q .

The final case for Condition 2 is $\bar{a} = a$. we find cases (23a), (23b), and (23g) apply yielding:

$$\begin{aligned} \sum_{z' \geq_A \bar{a}'} \nu^{a,\alpha}(\bar{a}, z') &= \frac{1}{|N|} \left(|N| - \sum_{z \in q \cup r} \hat{P}_{g(a,z)}^a(z_{g(a,z)}) \right. \\ &\quad \left. - \sum_{z' \in R} \hat{P}_{g(\alpha^T, z')}^a(1) \right) \\ &= \frac{1}{|N|} \sum_{z \in f(a)} \left(1 - \hat{P}_{g(a,z)}^a(z_{g(a,z)}) \right) \\ &= \frac{1}{|N|} \sum_{i \in N} \hat{P}_i^a(a_i) = \hat{P}^a(\bar{a}) \end{aligned} \quad (27)$$

where the first equality follows as sums over $Q \cup R$ are equivalent to the sums over Q and R as Q, R are disjoint, and that $z' \in R \Leftrightarrow z'_{g(\alpha^T, z')} = 1$ by definition of R . The second equality follows as the R sum is equivalent to one over s by bijection b^{a,α^T} , and then we may combine it with the sum over $q \cup r$, to a sum over $f(a)$ and $|f(a)| = |N|$. This concludes all cases for Condition 2. We omit arguments for Condition 3 as they run parallel to Condition 2.

To verify Condition 1, we consider each case of (23) separately. Equations (23b), (23d), (23e), (23f), and (23h) are trivial as these probabilities are well defined by definition. Lemma 2 provides:

$$\mathbb{P}_i^{\alpha}(1) \geq \hat{P}_i^a(1) \Leftrightarrow \hat{P}_i^a(0) \geq \mathbb{P}_i^{\alpha}(0) \quad (28)$$

where the right hand side follows from $\mathbb{P}_i(1, a', w) + \mathbb{P}_i(0, a', w) = 1 = \mathbb{P}_i(1, a', w_0) + \mathbb{P}_i(0, a', w_0)$. Equation (23a) follows directly from the hypothesis and (23c) holds from the right side of the equivalence.

The lone remaining case is (23g), for which we define sets $N_q = \{g(a, z) \mid z \in q\}$, $N_Q = \{g(\alpha^T, z) \mid z \in Q\}$ and so on for r, s, R, S . For convenience we denote unions of these sets as $N_{qr} := N_q \cup N_r$, $N_{QR} := N_Q \cup N_R$ and so on for other combinations of q, r, s and Q, R, S . Recalling q, r, Q, R are partitions over states that a, α^T may transition to, similarly, N_{qr}, N_{QR} are partitions of agents whose unilateral deviations result in such transitions. These sets are subsets of N and enable us to expand (23g)

$$\begin{aligned} \nu^{a,\alpha}(\bar{a}, \bar{a}') &= \frac{1}{|N|} \left(\sum_{i \in N_{qr} \cap N_{QR}} (1 - \hat{P}_i^a(-a_i) \right. \\ &\quad \left. - \mathbb{P}_i^{\alpha}(-\alpha_i^T)) \right. \\ &\quad \left. + \sum_{i \in N_{qr} \setminus N_{QR}} (1 - \hat{P}_i^a(-a_i)) \right. \\ &\quad \left. + \sum_{i \in N_{QR} \setminus N_{qr}} (1 - \mathbb{P}_i^{\alpha}(-\alpha_i^T)) \right) \end{aligned} \quad (29)$$

$$\nu^{a,\alpha}(\bar{a}, \bar{a}') = \begin{cases} \frac{1}{|N|} \left(\mathbb{P}_{g(\alpha^T, \bar{a}')}^\alpha(1) - \hat{\mathbb{P}}_{g(a', \bar{a}')}^a(1) \right) & \bar{a} = a, \bar{a}' \in R & (23a) \\ \frac{1}{|N|} \mathbb{P}_{g(\alpha^T, \bar{a}')}^\alpha(\bar{a}'_{g(\alpha^T, \bar{a}')}) & \bar{a} = a, \bar{a}' \in Q & (23b) \\ \frac{1}{|N|} \left(\hat{\mathbb{P}}_{g(a, \bar{a})}^a(0) - \mathbb{P}_{g(a, \bar{a})}^\alpha(0) \right) & \bar{a} \in r, \bar{a}' = \alpha^T & (23c) \\ \frac{1}{|N|} \hat{\mathbb{P}}_{g(a, \bar{a})}^a(\bar{a}_{g(a, \bar{a})}) & \bar{a} \in q, \alpha^T = \bar{a}' & (23d) \\ \frac{1}{|N|} \hat{\mathbb{P}}_{g(a, \bar{a})}^a(1) & \bar{a} = b^{a, \alpha^T}(\bar{a}'), \bar{a}' \in R & (23e) \\ \frac{1}{|N|} \mathbb{P}_{g(\alpha^T, \bar{a}')}^\alpha(0) & \bar{a} \in r, \bar{a}' = b^{a, \alpha^T}(\bar{a}) & (23f) \\ \frac{1}{|N|} \left(|N| - \sum_{z \in q \cup r} \hat{\mathbb{P}}_{g(a, z)}^a(z_{g(a, z)}) - \sum_{z' \in Q \cup R} \mathbb{P}_{g(\alpha^T, z')}^\alpha(z'_{g(\alpha^T, z')}) \right) & a = \bar{a}, \alpha^T = \bar{a}' & (23g) \\ 0 & \text{otherwise.} & (23h) \end{cases}$$

Fig. 2. The full specification of the one-step monotone coupling for Theorem 2. We adopt the notational convention that q, s, Q, S are assumed to take arguments a, a' and r, R take the argument a, a' respectively.

This expansion takes advantage of $|N| = |f(a)|$ which allows $|N|$ to enter the sums as 1. It now suffices to show that the summand of each sum is a well defined probability, then the sum of $|N|$ well defined probabilities divided by $|N|$ must also be a well defined probability. The last two terms in (29) are clearly well-defined probabilities, leaving only the first term.

We begin by investigating $i \in N_{qr} \cap N_{QR}$. In particular, we have $N_q = N_Q, N_s = N_R, N_r = N_S$ due to b^{a, α^T} and its bijectiveness due to Lemmas 3. By disjointness of q, r we have $N_{qr} = N_{QS}$ which we apply to $N_{qr} \cap N_{QR} = N_{QS} \cap N_{QR} = N_Q = N_q$. Applying definitions of q, Q we find $i \in N_q \implies \neg a_i = 1, \neg \alpha_i^T = 0$. Thus the summand of the first sum for $i \in N_q$ is given by

$$1 - \hat{\mathbb{P}}_i^a(1) - \mathbb{P}_i^\alpha(0) \geq 1 - \mathbb{P}_i^\alpha(1) - \mathbb{P}_i^\alpha(0) = 0 \quad (30)$$

where in the inequality is by (28), giving that the summands in the first term of (29) are themselves well defined probabilities. As all conditions have been met, $\nu^{a, \alpha}$ is a monotone coupling as desired. ■

D. A monotone coupling over histories

We now present coupling $\nu_{\pi}^{\hat{g}}$ which is constructed using the one-step coupling. Using this coupling we then go on to prove Lemma 1. Note, we define an indicator functions as $\mathbb{1}(a = a')$ to be 1 if $a = a'$ and 0 else for $a, a' \in A$.

Theorem 3: Let $g \in \mathcal{G}$ be an aligned history-dependent game and \hat{g} be its corresponding exact potential game. Then $\nu_{\pi}^{\hat{g}} : \mathcal{A}_T^2 \rightarrow [0, 1]$ is a monotone coupling between \hat{P}_{π}, P_{π} . This coupling is given by

$$\nu_{\pi}^{\hat{g}}(\alpha, \bar{\alpha}) = \pi(\alpha^1) \mathbb{1}(\alpha^1 = \bar{\alpha}^1) \prod_{t=1}^{T-1} \nu^{\alpha^t, \bar{\alpha}^{\leq t}}(\alpha^{t+1}, \bar{\alpha}^{t+1}) \quad (31)$$

where $\alpha, \bar{\alpha} \in \mathcal{A}_T, \pi \in \Delta(A)$.

Proof: Let $\alpha, \bar{\alpha} \in \mathcal{A}_T$ and let \hat{g} be a fixed corresponding fixed game to $g \in \mathcal{G}$. We begin by showing that if $\alpha \not\leq_{\mathcal{A}_T} \bar{\alpha}$, then $\nu_{\pi}^{\hat{g}}(\alpha, \bar{\alpha}) = 0$. Immediately, we have $\nu_{\pi}^{\hat{g}}(\alpha, \bar{\alpha}) = 0$ if $\alpha^1 \neq \bar{\alpha}^1$, so we need only consider cases where $\alpha^1 = \bar{\alpha}^1$. Inductively we find that if $\alpha \not\leq_{\mathcal{A}_T} \bar{\alpha}$ there

must exist some $t \in \{1, 2, 3, \dots, T-1\}$ such that $\alpha^t \leq_A \bar{\alpha}^t$ but $\alpha^{t+1} \not\leq_A \bar{\alpha}^{t+1}$, and let t be the minimal such value. In this case we have $\nu^{\alpha^t, \bar{\alpha}^{\leq t}}(\alpha^{t+1}, \bar{\alpha}^{t+1}) = 0$ because $\nu^{\alpha^t, \bar{\alpha}^{\leq t}}$ is a well defined monotone coupling by Theorem 2, yielding $\nu_{\pi}^{\hat{g}}(\alpha, \bar{\alpha}) = 0$ as desired. It also follows that $\nu_{\pi}^{\hat{g}}$ will always yield a well defined probability as it is either 0 or a product of well defined probabilities. Thus we only need to show that the marginal probabilities are preserved given by (15). We begin by showing the left equation of (15), that is:

$$\sum_{\alpha \leq_{\mathcal{A}_T} z} \nu_{\pi}^{\hat{g}}(\alpha, z) = \hat{P}_{\pi}(\alpha) \text{ for each } z \in \mathcal{A}_T \quad (32)$$

and omit the proof for the right hand equation as it proceeds identically. By inspecting (31), we only need to consider z such that $z^1 = \alpha^1$ and z features at most a single unilateral deviation between any $t, t+1$. With these two conditions we rewrite

$$\begin{aligned} \sum_{\alpha \leq_{\mathcal{A}_T} z} \nu_{\pi}^{\hat{g}}(\alpha, z) &= \sum_{\alpha \leq_{\mathcal{A}_T} z} \pi(\alpha^1) \prod_{t=1}^{T-1} \nu^{\alpha^t, z^{\leq t}}(\alpha^{t+1}, z^{t+1}) \\ &= \pi(\alpha^1) \sum_{\alpha^2 \leq_A z^2} \nu^{\alpha^1, z^{\leq 1}}(\alpha^2, z^2) \dots \\ &\quad \sum_{\alpha^T \leq_{\mathcal{A}_T} z^T} \nu^{\alpha^{T-1}, z^{\leq T-1}}(\alpha^T, z^T). \end{aligned} \quad (33)$$

as the combinatorial form. Critically, this allows us to to apply the marginal sum properties of $\nu^{\alpha^t, z^{\leq t}}$ from Theorem 2 for each $t \in \{1, 2, \dots, T\}$. First, considering the rightmost sum in (33), it holds that

$$\sum_{\alpha^T \leq_{\mathcal{A}_T} z^T} \nu^{\alpha^{T-1}, z^{\leq T-1}}(\alpha^T, z^T) = \hat{P}^{\alpha^{T-1}}(\alpha^T). \quad (34)$$

Because this has no dependence on z we may factor out $\hat{P}^{\alpha^{T-1}}(\alpha^T)$ and repeat the process on the new rightmost sum. After performing this process recursively on all sums, we have

$$\sum_{\alpha \leq_{\mathcal{A}_T} z} \nu_{\pi}^{\hat{g}}(\alpha, z) = \pi(\alpha^1) \prod_{t=1}^{T-1} \hat{P}^{\alpha^t}(\alpha^{t+1}) = \hat{P}_{\pi}(\alpha) \quad (35)$$

as desired, noting we accounted for the indicator functions in $\nu_{\pi}^{\hat{g}}$. This concludes the proof of Theorem 3. ■

Now that the necessary results have been developed we proceed with the proof of Lemma 1.

Proof of Lemma 1: Let $g \in \mathcal{G}^A$ be an aligned history-dependent game and $\mathcal{I} \subset \mathcal{A}_T$ be an upper set. Define $\mathbb{1}_{\mathcal{I}}(a) := \mathbb{1}(a \in \mathcal{I})$ as an indicator function. Consider probability measures P_{π}, \hat{P}_{π} coupled by $\nu_{\pi}^{\hat{g}}$ in Theorem 3, we have

$$\begin{aligned} P_{\pi}(\mathcal{I}) - \hat{P}_{\pi}(\mathcal{I}) &= \mathbb{E}_{P_{\pi}}(\mathbb{1}_{\mathcal{I}}) - \mathbb{E}_{\hat{P}_{\pi}}(\mathbb{1}_{\mathcal{I}}) \\ &= \nu_{\pi}^{\hat{g}}(\mathcal{I}^c, \mathcal{I}) \geq 0. \end{aligned} \quad (36)$$

where the second equality follows by Proposition 3 as $\mathbb{1}_{\mathcal{I}}$ is increasing in \mathcal{A}_T . Note (36) runs parallel to the proof of [24, Corollary 3]. That is, for any upper set $\mathcal{I} \subset \mathcal{A}_T$ we have

$$P_{\pi}(\mathcal{I}) \geq \hat{P}_{\pi}(\mathcal{I}) \quad (37)$$

meaning we have stochastic dominance. Let $((\vec{0})_{t=1}^{T-1}, \vec{1}) \in \mathcal{I}$. This induces $\bar{\mathcal{I}}$ such that it includes every path such that at time T the $\vec{1}$ state is played. This yields the following interpretation

$$P_{\pi}(\bar{\mathcal{I}}) = \Pr(s(T; \tau, \pi, g) = \vec{1}) \quad (38)$$

representing the probability that at time T game g is in the $\vec{1}$ action profile given initial distribution $\pi \in \Delta(A)$ and learning temperature parameter τ . Noting a parallel interpretation to (38) holds for \hat{P}_{π}, \hat{g} , we apply these to (37) to obtain

$$\Pr(s(T; \tau, \pi, g) = \vec{1}) \geq \Pr(s(T; \tau, \pi, \hat{g}) = \vec{1}) \quad (39)$$

as desired. ■

REFERENCES

- [1] J. Howard, A. Huang, Z. Li, Z. Tufekci, V. Zdimal, H.-M. van der Westhuizen, A. von Delft, A. Price, L. Fridman, L.-H. Tang, V. Tang, G. L. Watson, C. E. Bax, R. Shaikh, F. Questier, D. Hernandez, L. F. Chu, C. M. Ramirez, and A. W. Rimoim, "An evidence review of face masks against covid-19," *Proceedings of the National Academy of Sciences*, vol. 118, no. 4, 2021.
- [2] L. Matrajt and T. Leung, "Evaluating the effectiveness of social distancing interventions to delay or flatten the epidemic curve of coronavirus disease," *Emerging infectious diseases*, vol. 26, no. 8, p. 1740, 2020.
- [3] H. P. Young, "The evolution of conventions," *Econometrica: Journal of the Econometric Society*, pp. 57–84, 1993.
- [4] M. Kearns, M. L. Littman, and S. Singh, "Graphical models for game theory," *arXiv preprint arXiv:1301.2281*, 2013.
- [5] J. R. Marden and J. S. Shamma, "Revisiting log-linear learning: Asynchrony, completeness and payoff-based implementation," *Games and Economic Behavior*, vol. 75, no. 2, pp. 788–808, 2012.
- [6] R. Chandan, D. Paccagnan, and J. R. Marden, "When Smoothness is Not Enough: Toward Exact Quantification and Optimization of the Price-of-Anarchy," in *58th IEEE Conference on Decision and Control*, pp. 4041–4046, 2019.
- [7] K. Paarporn, B. Canty, P. N. Brown, M. Alizadeh, and J. R. Marden, "The Impact of Complex and Informed Adversarial Behavior in Graphical Coordination Games," *IEEE Transactions on Control of Network Systems*, vol. 8, no. 1, pp. 200–211, 2020.
- [8] J. R. Marden and A. Wierman, "Distributed welfare games," *Operations Research*, vol. 61, no. 1, pp. 155–168, 2013.
- [9] A. Kanakia, B. Touri, and N. Correll, "Modeling multi-robot task allocation with limited information as global game," *Swarm Intelligence*, vol. 10, no. 2, pp. 147–160, 2016.
- [10] C. Wang, C. Xu, X. Yao, and D. Tao, "Evolutionary generative adversarial networks," *IEEE Transactions on Evolutionary Computation*, vol. 23, no. 6, pp. 921–934, 2019.
- [11] U. Garciarena, R. Santana, and A. Mendiburu, "Evolved gans for generating pareto set approximations," in *Proceedings of the Genetic and Evolutionary Computation Conference*, pp. 434–441, 2018.

- [12] V. Costa, N. Lourenço, J. Correia, and P. Machado, "Coegan: evaluating the coevolution effect in generative adversarial networks," in *Proceedings of the Genetic and Evolutionary Computation Conference*, pp. 374–382, 2019.
- [13] A. R. Tilman, J. R. Watson, and S. Levin, "Maintaining cooperation in social-ecological systems," *Theoretical Ecology*, vol. 10, no. 2, pp. 155–165, 2017.
- [14] A. R. Tilman, J. B. Plotkin, and E. Akçay, "Evolutionary games with environmental feedbacks," *Nature communications*, vol. 11, no. 1, pp. 1–11, 2020.
- [15] S. Skoulakis, T. Fiez, R. Sim, G. Piliouras, and L. Ratliff, "Evolutionary game theory squared: Evolving agents in endogenously evolving zero-sum games," *arXiv preprint arXiv:2012.08382*, 2020.
- [16] J. S. Weitz, C. Eksin, K. Paarporn, S. P. Brown, and W. C. Ratcliff, "An oscillating tragedy of the commons in replicator dynamics with game-environment feedback," *Proceedings of the National Academy of Sciences*, vol. 113, no. 47, pp. E7518–E7525, 2016.
- [17] J. A. Weill, M. Stigler, O. Deschenes, and M. R. Springborn, "Social distancing responses to covid-19 emergency declarations strongly differentiated by income," *Proceedings of the National Academy of Sciences*, vol. 117, no. 33, pp. 19658–19660, 2020.
- [18] B. Jeffrey, C. E. Walters, K. E. Ainslie, O. Eales, C. Ciavarella, S. Bhatia, S. Hayes, M. Baguelin, A. Boonyasiri, N. F. Brazeau, *et al.*, "Anonymised and aggregated crowd level mobility data from mobile phones suggests that initial compliance with covid-19 social distancing interventions was high and geographically consistent across the uk," *Wellcome Open Research*, vol. 5, 2020.
- [19] J. Jay, J. Bor, E. O. Nsoesie, S. K. Lipson, D. K. Jones, S. Galea, and J. Raifman, "Neighbourhood income and physical distancing during the covid-19 pandemic in the united states," *Nature human behaviour*, vol. 4, no. 12, pp. 1294–1302, 2020.
- [20] C. Alós-Ferrer and N. Netzer, "The logit-response dynamics," *Games and Economic Behavior*, vol. 68, no. 2, pp. 413–427, 2010.
- [21] P. N. Brown, H. P. Borowski, and J. R. Marden, "Security against impersonation attacks in distributed systems," *IEEE Transactions on Control of Network Systems*, vol. 6, no. 1, pp. 440–450, 2019.
- [22] B. S. Pradelski and H. P. Young, "Learning efficient nash equilibria in distributed systems," *Games and Economic Behavior*, vol. 75, no. 2, pp. 882–897, 2012.
- [23] C. Nowzari, V. M. Preciado, and G. J. Pappas, "Analysis and Control of Epidemics: A Survey of Spreading Processes on Complex Networks," *IEEE Control Systems*, vol. 36, no. 1, pp. 26–46, 2016.
- [24] K. Paarporn, C. Eksin, J. S. Weitz, and J. S. Shamma, "Networked SIS Epidemics with Awareness," *IEEE Transactions on Computational Social Systems*, vol. 4, no. 3, pp. 93–103, 2017.

APPENDIX

Proof of Lemma 3: Let $a, a' \in A$ such that $a' \geq_A a$. We proceed by proving $b^{a,a'} : r(a) \rightarrow S(a')$ is a bijection; the other bijection statements are proved similarly.

We begin by proving injectiveness, that is $b^{a,a'}(z) = b^{a,a'}(z') \implies z = z'$ for $z, z' \in r(a)$. Observe $g(a, z) = g(a', b^{a,a'}(z)) = g(a', b^{a,a'}(z')) = g(a, z')$ where the first and third inequalities follow by definition of $b^{a,a'}$ and the middle by hypothesis. Injectiveness follows from $g(a, z) = g(a, z')$ meaning a, z and a, z' differ by the same agent's unilateral deviation. In that context, the possible actions agent $g(a, z)$ is given by $A_{g(a,z)} \setminus \{a_{g(a,z)}\}$ which is a singleton by the binary action property, leaving only one possible state a could transition to in $r(a)$ via a unilateral deviation. Thus $z = z'$ as desired.

Next we show surjection, that is for any $z' \in S(a, a')$ there exists a $z \in r(a)$ such that $b^{a,a'}(z) = z'$, for $a, a' \in A$ and $a \leq_A a'$. By definition of $S(a, a')$, $z' \not\geq_A a'$, but as $z' \in f(a')$ z', a' differ by only a single unilateral deviation by some agent i . By partial ordering \leq_A we may infer $a'_i = 1, z'_i = 0$ else $z' \not\geq_A a'$ would be violated. Further, we may infer $a = 1$ as suppose $a = 0$, then $z' \in Q(a, a')$, giving a contradiction

to the definition of z' . It is easy to see by definition of $r(a)$ that $a_i = 1 \implies z \in r(a)$ satisfying $g(a, z) = g(a', z')$ as $z_i \neq a_i$ but $z_{-i} = a_{-i}$ by $z \in f(a)$. Note $g(a, z) = g(a', z')$ is always satisfied when $b^{a, a'}(z) = z'$ by definition of the function. ■